



Problems based on the interval $(-l, l)$

Even function:

A function is said to be even if $f(x) = f(-x)$. i.e., If $f(x)$ is an even function,

$$\int_{-l}^l f(x) dx = 2 \int_0^l f(x) dx$$

Odd function:

A function is said to be odd if

$$f(x) = -f(-x) \text{ or } f(-x) = -f(x). \text{ If}$$

$f(x)$ is an odd function,

$$\int_{-l}^l f(x) dx = 0.$$

Note:

If $f(x)$ does not satisfy even and odd function then it is called neither even nor odd function.

Example:

1. $f(x) = x^2$ in $(-\pi, \pi)$ - Even
2. $f(x) = x \cos x$ in $(-l, l)$ - Odd
3. $f(x) = x \sin x$ in $(-\pi, \pi)$ - Even
4. $f(x) = |x|$ in $(-2, 2)$ - Even
5. $f(x) = (l-x)^2$ in $(-l, l)$ - Neither even nor odd
6. $f(x) = x + x^2$ in $(-\pi, \pi)$ - Neither even nor odd
7. $f(x) = x - x^2$ in $(-2, 2)$ - Neither even nor odd



Note:

For split up problem,

$$f(x) = \begin{cases} \varphi_1(x) \\ \varphi_2(x) \end{cases}$$

Even $\rightarrow \varphi_1(x) = \varphi_2(-x)$

Odd $\rightarrow \varphi_1(x) = -\varphi_2(-x)$.

Example:

* $f(x) = \begin{cases} L+x, & -L < x < 0 \\ L-x, & 0 < x < L \end{cases}$

$\varphi_1(x) = L+x, \varphi_1(-x) = L-x = \varphi_2(x)$

$\varphi_2(x) = L-x, \varphi_2(-x) = L+x = \varphi_1(x)$

$\therefore f(x)$ is an even function.

* $f(x) = \begin{cases} -k, & -\pi \leq x \leq 0 \\ k, & 0 \leq x \leq \pi \end{cases}$

$\varphi_1(x) = -k, \varphi_1(-x) = -k = -\varphi_2(x)$

$\varphi_2(x) = k, \varphi_2(-x) = k = -\varphi_1(x)$

$\therefore f(x)$ is an odd function.

Note:

* Even \times Even = Even

* Odd \times Odd = Even

* Even \times Odd = Odd

* Odd \times Even = Odd.



Interval $(-l, l)$		
$f(x)$ is an even function	$f(x)$ is an odd function	$f(x)$ is neither even nor odd
$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \frac{\cos \frac{n\pi x}{l}}{l} + \sum_{n=1}^{\infty} b_n \frac{\sin \frac{n\pi x}{l}}{l}$		
$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx$ $= \frac{2}{l} \int_0^l f(x) dx$	$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx$ $a_0 = 0$	$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx$
$a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$ $= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$	$a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$ $a_n = 0$	$a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$
$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx$ $b_n = 0$	$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \left(\frac{n\pi x}{l} \right) dx$ $b_n = \frac{2}{l} \int_0^l f(x) \sin \left(\frac{n\pi x}{l} \right) dx$	$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \left(\frac{n\pi x}{l} \right) dx$



① Find the Fourier series for the function

$$f(x) = \begin{cases} L+x, & (-L, 0) \\ L-x, & (0, L) \end{cases}$$

Soln:

Here $l = L$

$$\phi_1(x) = L+x$$

$$\phi_1(-x) = L-x = \phi_2(x)$$

$$\phi_2(x) = L-x$$

$$\phi_2(-x) = L-(-x) = L+x = \phi_1(x)$$

$\therefore f(x)$ is an even function.

$$b_n = 0$$

Fourier Series :

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{L}\right)$$

To find a_0 :

$$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx$$

$$= \frac{2}{L} \int_0^L f(x) dx$$

$$= \frac{2}{L} \int_0^L (L-x) dx$$

$$= \frac{2}{L} \left[Lx - \frac{x^2}{2} \right]_0^L$$

$$a_0 = L$$



To find a_n :

$$\begin{aligned}
 a_n &= \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx \\
 &= \frac{2}{L} \int_0^L (L-x) \cos \frac{n\pi x}{L} dx \\
 &= \frac{2}{L} \left[\frac{L}{n\pi} (L-x) \sin \left(\frac{n\pi x}{L} \right) - \frac{L^2}{n^2 \pi^2} \cos \left(\frac{n\pi x}{L} \right) \right]_0^L \\
 &= \frac{2}{L} \left[-\frac{L^2}{n^2 \pi^2} \cos n\pi + \frac{L^2}{n^2 \pi^2} \cos 0 \right] \\
 &= \frac{2}{L} \cdot \frac{L^2}{n^2 \pi^2} [1 - (-1)^n]
 \end{aligned}$$

$$a_n = \frac{2L}{n^2 \pi^2} [1 - (-1)^n]$$

Subs a_0 , a_n and b_n in (1),

$$f(x) = \frac{L}{2} + \sum_{n=1}^{\infty} \frac{2L}{n^2 \pi^2} [1 - (-1)^n] \cos \left(\frac{n\pi x}{L} \right)$$

2) $f(x) = |x|$ in $-l \leq x \leq l$.

Soln:

$f(x)$ is an even function

$$b_n = 0.$$

$$a_0 = l$$

$$a_n = \frac{2l}{n^2 \pi^2} [(-1)^n - 1]$$

3) $f(x) = |x|$ in $-\pi \leq x \leq \pi$

Soln:



$f(x)$ is an even function

$$b_n = 0$$

$$a_0 = \pi$$

$$a_n = \frac{2}{\pi n^2} [(-1)^n - 1]$$

$$\textcircled{4} \quad f(x) = \begin{cases} 1+x, & -2 \leq x \leq 0 \\ 1-x, & 0 \leq x \leq 2 \end{cases}$$

Soln:

$f(x)$ is an even fn.

$$b_n = 0$$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{2} \rightarrow \textcircled{1}$$

To find a_0 :

$$\begin{aligned} a_0 &= \frac{1}{l} \int_{-l}^l f(x) dx \\ &= \frac{2}{l} \int_0^l f(x) dx \\ &= \frac{2}{2} \int_0^2 (1-x) dx \end{aligned}$$

$$a_0 = 0$$

To find a_n :

$$a_n = \int_0^2 (1-x) \cos \left(\frac{n\pi x}{2} \right) dx$$

$$a_n = \frac{4}{\pi^2 n^2} [1 - (-1)^n]$$

$$\therefore f(x) = \sum_{n=1}^{\infty} \frac{4}{\pi n^2} [1 - (-1)^n] \cos \left(\frac{n\pi x}{2} \right)$$



5) $f(x) = x - x^2, -l \leq x \leq l.$

Soln:

$f(x)$ is neither even nor odd function.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

To find a_0 :

$$a_0 = \frac{1}{l} \int_{-l}^l (x - x^2) dx$$

$$a_0 = -\frac{2l^2}{3}$$

To find a_n :

$$a_n = \frac{1}{l} \int_{-l}^l (x - x^2) \cos \frac{n\pi x}{l} dx$$

$$a_n = \frac{-4l^2(-1)^n}{n^2\pi^2}$$

To find b_n :

$$b_n = \frac{1}{l} \int_{-l}^l (x - x^2) \sin \frac{n\pi x}{l} dx$$

$$b_n = \frac{-2l(-1)^n}{n\pi}$$

Subs a_0, a_n & b_n in ①,

$$f(x) = -\frac{l^2}{3} + \sum_{n=1}^{\infty} \frac{-4l^2(-1)^n}{n^2\pi^2} \cos\left(\frac{n\pi x}{l}\right) + \sum_{n=1}^{\infty} \frac{-2l(-1)^n}{n\pi} \sin\left(\frac{n\pi x}{l}\right).$$