



Continuous Random Variable :

A random variable 'x' is called a continuous random variable if it takes all possible values in a given interval.

Examples : Age , Height and Weight

Distribution function (or) Cumulative Distribution function of the random variable X :

The c.d.f of a continuous random variable x is defined as ,

$$F(x) = P(X \leq x) = \int_{-\infty}^x f(x) dx$$

Probability Density function: (P.D.f)

Let x be a continuous random variable. The function f(x) is called the p.d.f of the random variable x if it satisfies the following conditions :

(i) $f(x) \geq 0$, $-\infty < x < \infty$

(ii) $\int_{-\infty}^{\infty} f(x) dx = 1$

Remark :

1. $P(a < x < b) = P(a \leq x \leq b) = \int_a^b f(x) dx$

2. $P(x > a) = \int_a^{\infty} f(x) dx$

3. $P(x < a) = \int_{-\infty}^a f(x) dx$

4. $P(x > a | x > b) = \frac{P(x > a)}{P(x > b)}$



DEPARTMENT OF MATHEMATICS

④ If 'x' is a Continuous random variable whose p.d.f is given by,

$$f(x) = \begin{cases} c(4x - 2x^2), & 0 < x < 2 \\ 0, & \text{otherwise} \end{cases}$$

Find (a) what is the value of 'c'? (b) Find $P(x > 1)$

Solution:

(a) Given: $f(x) = \begin{cases} c(4x - 2x^2), & 0 < x < 2 \\ 0, & \text{otherwise} \end{cases} \longrightarrow \textcircled{1}$

$$\int_{-\infty}^{\infty} f(x) dx = 1$$

$$\therefore \int_0^2 c(4x - 2x^2) dx = 1$$

$$c \left[4 \frac{x^2}{2} - 2 \frac{x^3}{3} \right]_0^2 = 1$$

$$c \left[2(2^2) - \frac{2}{3}(2^3) \right] = 1$$

$$c \left[8 - \frac{16}{3} \right] = 1 \Rightarrow c \left(\frac{24 - 16}{3} \right) = 1$$

$$c \left(\frac{8}{3} \right) = 1$$

$$\boxed{c = \frac{3}{8}}$$

Put $c = \frac{3}{8}$ in $\textcircled{1}$,

$$f(x) = \begin{cases} \frac{3}{8}(4x - 2x^2), & 0 < x < 2 \\ 0, & \text{otherwise} \end{cases}$$



$$\begin{aligned} \text{(b) } P(x > 1) &= \int_1^{\infty} f(x) dx \\ &= \int_1^2 f(x) dx + \int_2^{\infty} f(x) dx \\ &= \int_1^2 \frac{3}{8} (4x - 2x^2) dx + 0 \\ &= \frac{3}{8} \times 2 \int_1^2 (2x - x^2) dx \\ &= \frac{3}{4} \left[\frac{2x^2}{2} - \frac{x^3}{3} \right]_1^2 \\ &= \frac{3}{4} \left[(4 - 1) - \frac{1}{3} (8 - 1) \right] \\ &= \frac{3}{4} \left[3 - \frac{7}{3} \right] = \frac{3}{4} \left[\frac{9-7}{3} \right] = \frac{2}{4} = \frac{1}{2} \end{aligned}$$

$$P(x > 1) = \frac{1}{2}$$

⑤ The amount of time, in hours, that a computer functions before breaking down is a continuous random variable with probability density function given by,

$$f(x) = \begin{cases} \lambda e^{-x/100}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

What is the probability that (a) a computer will function between 50 and 150 hrs, before breaking down (b) it will function less than 500 hours.



Solution:

$$\text{Given: } f(x) = \begin{cases} \lambda e^{-x/100} & , x \geq 0 \\ 0 & , x < 0 \end{cases} \longrightarrow \textcircled{1}$$

Since $f(x)$ is a p.d.f of 'x',

$$\int_{-\infty}^{\infty} f(x) dx = 1$$

$$\int_{-\infty}^0 f(x) dx + \int_0^{\infty} f(x) dx = 1$$

$$0 + \int_0^{\infty} \lambda e^{-x/100} dx = 1$$

$$\lambda \left[\frac{e^{-x/100}}{-1/100} \right]_0^{\infty} = 1$$

$$\lambda (-100) [e^{-\infty} - e^0] = 1$$

$$\lambda (-100) [0 - 1] = 1$$

$$100\lambda = 1$$

$$\boxed{\lambda = \frac{1}{100}}$$

(a) We know that,

$$P(a < x < b) = \int_a^b f(x) dx$$

$$P(50 < x < 150) = \int_{50}^{150} f(x) dx$$

$$= \int_{50}^{150} \frac{1}{100} e^{-x/100} dx$$



DEPARTMENT OF MATHEMATICS

$$\begin{aligned} &= \frac{1}{100} \left[\frac{e^{-x/100}}{-1/100} \right]_{50}^{150} \\ &= - \left[e^{-150/100} - e^{-50/100} \right] \\ &= - e^{-1.5} + e^{-0.5} \\ &= - 0.2231 + 0.6065 \\ &= \underline{0.3834} \end{aligned}$$

$$\begin{aligned} \text{(b) } P(x < 500) &= \int_0^{500} f(x) dx \\ &= \int_0^{500} \frac{1}{100} e^{-x/100} dx \\ &= \frac{1}{100} \left[\frac{e^{-x/100}}{-1/100} \right]_0^{500} \\ &= - \left[e^{-500/100} - e^0 \right] \\ &= - \left[e^{-5} - 1 \right] \\ &= 1 - e^{-5} \\ &= 1 - 0.0067 \end{aligned}$$

$$\boxed{P(x < 500) = 0.9933}$$