



SNS COLLEGE OF ALLIED HEALTH SCIENCES
SNS Kalvi Nagar, Coimbatore - 35
Affiliated to Dr MGR Medical University, Chennai



DEPARTMENT OF CARDIO PULMONARY PERFUSION CARE
TECHNOLOGY

COURSE NAME : Pharmacology Pathology and Clinical Microbiology

2nd YEAR


TOPIC : COAGULANTS

Coagulants

Coagulant

- Haemostasis (arrest of blood loss) and blood coagulation involve complex interactions between the injured vessel wall, platelets and coagulation factors.

Mechanisms of blood coagulation




Coagulants

Vitamin K	K1 (from plants fat-soluble):	Phytonadione (Phylloquinone)
	K3 (synthetic)	
	—Fat-soluble:	Menadione, Acetomenaphthone
	—Water-soluble:	Menadione sod. Bisulfite, Menadione, sod. Diphosphate
Miscellaneous	Fibrinogen (human), Antihæmophilic factor, Desmopressin, Adrenochrome monosemicarbazone, Rutin, Ethamsylate	


Vitamin K

- Vit. K is a fat-soluble dietary principle required for the synthesis of clotting factors.
- **Daily requirement:** Vit. K₂ produced by colonic bacteria and 3–10 µg/day external source may be sufficient. The total requirement of Vit. K for an adult has been estimated to be 50–100 µg/day.

Vitamin K



Inactive form Functional form



Vit. K

Vit. K

Vitamin K

Action:

Vit K acts as a cofactor at a late stage in the synthesis by liver of coagulation proteins - prothrombin, factors VII, IX and X.

Use: The only use of vit K is in prophylaxis and treatment of bleeding due to deficiency of clotting factors.

Coagulants

Plasma fractions

- Deficiencies in plasma coagulation factors can cause bleeding.
- **Factor VIII deficiency (classic hemophilia or hemophilia A) and factor IX deficiency (Christmas disease, or hemophilia B)** account for most of the heritable coagulation defects. Concentrated plasma fractions and recombinant protein preparations are available for the treatment of these deficiencies.

Coagulants

Desmopressin acetate

- Desmopressin (DDAVP) stimulates the release of *von Willebrand factor (vWF)* from the **Weibel–Palade bodies** of **endothelial cells**, thereby increasing the levels of vWF (as well as coagulant factor VIII) 3 to 5-fold.
- It also used to promote the release of von *Willebrand factor* in patients with **coagulation disorders** such as *von Willebrand* disease, **mild hemophilia A** and **thrombocytopenia**.

Coagulants

Cryoprecipitate

- Cryoprecipitate is a plasma protein fraction obtainable from whole blood. It is used to treat deficiencies or qualitative abnormalities of fibrinogen.
- It may also be used for patients with factor VIII deficiency and von Willebrand disease

THANK YOU