

SNS COLLEGE OF ALLIED HEALTH SCIENCES SNS Kalvi Nagar, Coimbatore - 35 Affiliated to Dr MGR Medical University, Chennai

DEPARTMENT OF PHYSICIAN ASSISSTANT

COURSE NAME: PULMONOLOGY

TOPIC – BASICS OF RESPIRATORY MECHANISMS

B.SHANMUGA PRIYA Lecturer **OTAT SNSCAHS**





FUNCTIONS OF RESPIRATORY SYSTEM



- **Oxygen supplier.** The job of the respiratory system is to keep the body constantly supplied with oxygen.
- **Elimination.** Elimination of carbon dioxide.
- **Gas exchange.** The respiratory system organs oversee the gas exchanges that occur between the **blood** and the external environment.
- **Passageway.** Passageways that allow air to reach the lungs.
- **Humidifier.** Purify, humidify, and warm incoming air.



The functions of the respiratory system are:



Anatomy of the Respiratory System

The organs of the respiratory system include the nose, pharynx, larynx, trachea, bronchi, and their smaller branches, and the lungs, which contain the alveoli.







NOSE ANATOMY



The nose is the only externally visible part of the respiratory system

- **Nostrils.** During breathing, air enters the nose by \bullet passing through the nostrils, or nares.
- Nasal cavity. The interior of the nose consists of ulletthe nasal cavity, divided by a midline **nasal** septum.
- ullet
- ullet



Olfactory receptors. The olfactory receptors for the sense of <u>smell</u> are located in the mucosa in the slitlike superior part of the nasal cavity, just beneath the ethmoid bone.

Respiratory mucosa. The rest of the mucosal lining, the nasal cavity called the respiratory mucosa, rests on a rich network of thin-walled veins that warms the air as it flows past.



NOSE ANATOMY

- **Mucus.** In addition, the sticky mucus produced by the mucosa's glands moistens the air and traps ● incoming bacteria and other foreign debris, and **lysozyme enzymes** in the mucus destroy bacteria chemically.
- **Ciliated cells.** The ciliated cells of the nasal mucosa create a gentle current that moves the sheet of \bullet contaminated mucus posteriorly toward the throat, where it is swallowed and digested by stomach juices.
- **Conchae.** The <u>lateral</u> walls of the nasal cavity are uneven owing to three mucosa-covered projections, \bullet or lobes called conchae, which greatly increase the surface area of the mucosa exposed to the air, and also increase the air turbulence in the nasal cavity.
- **Palate.** The nasal cavity is separated from the oral cavity below by a partition, the palate; anteriorly, ulletwhere the palate is supported by bone, is the **hard palate**; the unsupported posterior part is the **soft** palate.
- **Paranasal sinuses.** The nasal cavity is surrounded by a ring of paranasal sinuses located in the frontal, • sphenoid, ethmoid, and maxillary <u>bones</u>; theses sinuses lighten the skull, and they act as a resonance chamber for speech.







Pharynx

- Size. The pharynx is a muscular passageway about 13 cm (5 inches) long that vaguely resembles a short length of red garden hose.
- Function. Commonly called the throat, the pharynx serves as a common passageway for food and air.
- **Portions of the pharynx.** Air enters the superior portion, the **nasopharynx**, from the nasal cavity and then descends through the **oropharynx** and **laryngopharynx** to enter the larynx below.
- **Pharyngotympanic tube.** The pharyngotympanic tubes, which drain the middle ear open into the nasopharynx.
- **Pharyngeal tonsil.** The pharyngeal tonsil, often called **adenoid** is located high in the nasopharynx.
- **Palatine** <u>tonsils</u>. The palatine <u>tonsils</u> are in the oropharynx at the end of the soft palate.
- Lingual <u>tonsils</u>. The lingual <u>tonsils</u> lie at the base of the tongue.





Larynx

- The larynx or **voice box** routes air and food into the proper channels and plays a role in speech.
- Structure. Located inferior to the pharynx, it is formed by eight rigid hyaline cartilages and a spoonshaped flap of elastic cartilage, the epiglottis.
- **Thyroid cartilage.** The largest of the hyaline cartilages is the shield-shaped thyroid cartilage, which protrudes anteriorly and is commonly called **Adam's apple**.
- Epiglottis. Sometimes referred to as the "guardian of the airways", the epiglottis protects the superior opening of the larynx.
- Vocal folds. Part of the mucous membrane of the larynx forms a pair of folds, called the vocal folds, or true vocal cords, which vibrate with expelled air and allows us to speak.
- **Glottis.** The slitlike passageway between the vocal folds is the glottis.



els and plays a role in speech. id hyaline cartilages and a spoon-





TRACHEA

- Length. Air entering the trachea or windpipe from the larynx travels down its length (10 to 12 cm or about 4 inches) to the level of the fifth thoracic vertebra, which is approximately midchest.
- **Structure.** The trachea is fairly rigid because its walls are reinforced with **C-shaped rings** of hyaline cartilage; the open parts of the rings abut the esophagus and allow it to expand anteriorly when we swallow a large piece of food, while the solid portions support the trachea walls and keep it patent, or open, in spite of the pressure changes that occur during breathing.
- **Cilia.** The trachea is lined with ciliated mucosa that beat continuously and in a direction opposite to that of the incoming air as they propel mucus, loaded with dust particles and other debris away from the lungs to the throat, where it can be swallowed or spat out





Main Bronchi

- **Structure.** The right and left main (primary) bronchi are formed by the division of the trachea.
- Location. Each main bronchus runs obliquely before it plunges into the medial <u>depression</u> of the lung on its own side.
- Size. The right main bronchus is wider, shorter, and straighter than the left



LUNGS







- Location. The lungs occupy the entire thoracic cavity except for the most central area, the mediastinum, which houses the heart, the great blood vessels, bronchi, esophagus, and other organs.
- **Apex.** The narrow, superior portion of each lung, the apex, is just deep into the clavicle.
- **Base.** The broad lung area resting on the diaphragm is the base.



LUNGS

- **Division.** Each lung is divided into lobes by fissures; the left lung has **two lobes**, and the right lung has **three**.
- Pleura. The surface of each lung is covered with a visceral serosa called the pulmonary, or visceral pleura, and the walls of the thoracic cavity are lined by the **parietal pleura**.
- **Pleural fluid.** The pleural membranes produce pleural fluid, a slippery serous secretion that allows the lungs to glide easily over the thorax wall during breathing movements and causes the two pleural layers to cling together.
- **Pleural space.** The lungs are held tightly to the thorax wall, and the pleural space is more of a potential space • than an actual one.
- **<u>Bronchioles</u>**. The smallest of the conducting passageways are the bronchioles. \bullet
- Alveoli. The terminal bronchioles lead to the respiratory zone structures, even smaller conduits that eventually terminate in alveoli or air sacs.
- **Respiratory zone.** The respiratory zone, which includes the respiratory bronchioles, alveolar ducts, alveolar sacs, and alveoli, is the only site of <u>gas exchange</u>.
- **Conducting zone structures.** All other respiratory passages are conducting zone structures that serve as \bullet conduits to and from the respiratory zone.
- **Stroma.** The balance of the lung tissue, its stroma, is mainly elastic connective tissue that allows the lungs to ulletrecoil passively as we exhale





The Respiratory Membrane

- **Wall structure.** The walls of the alveoli are composed largely of a single, thin layer of squamous epithelial cells.
- Alveolar pores. Alveolar pores connect neighboring air sacs and provide alternative routes for air to reach alveoli whose feeder bronchioles have been clogged by mucus or otherwise blocked.
- **Respiratory membrane.** Together, the alveolar and capillary walls, their fused basement membranes, \bullet and occasional elastic fibers construct the respiratory membrane (air-blood barrier), which has gas (air) flowing past on one side and blood flowing past on the other.
- **Alveolar macrophages.** Remarkably efficient alveolar macrophages sometimes called "dust cells", \bullet wander in and out of the alveoli picking up bacteria, carbon particles, and other debris.
- **Cuboidal cells.** Also scattered amid the epithelial cells that form most of the alveolar walls are chunky \bullet cuboidal cells, which produce a lipid (fat) molecule called **surfactant**, which coats the gas-exposed alveolar surfaces and is very important in lung function.





Physiology of the Respiratory System

The major function of the respiratory system is to supply the body with oxygen and to dispose of carbon dioxide. To do this, at least four distinct events, collectively called respiration, must occur.

Respiration

- **Pulmonary** <u>ventilation</u>. Air must move into and out of the lungs so that gasses in the air sacs are continuously refreshed, and this process is commonly called breathing.
- **External respiration.** Gas exchange between the pulmonary blood and alveoli must take ulletplace.
- **Respiratory gas transport.** Oxygen and carbon dioxide must be transported to and from ulletthe lungs and tissue cells of the body via the bloodstream.
- **Internal respiration.** At systemic capillaries, gas exchanges must be made between the \bullet blood and tissue cells.





Mechanics of Breathing

- **Rule.** Volume changes lead to pressure changes, which lead to the flow of gasses to equalize pressure. \bullet
- **Inspiration.** Air is flowing into the lungs; the chest is expanded laterally, the rib cage is elevated, and \bullet the diaphragm is depressed and flattened; lungs are stretched to the larger thoracic volume, causing the intrapulmonary pressure to fall and air to flow into the lungs.
- **Expiration.** Air is leaving the lungs; the chest is depressed and the lateral dimension is reduced, the rib \bullet cage is descended, and the diaphragm is elevated and dome-shaped; lungs recoil to a smaller volume, intrapulmonary pressure rises, and air flows out of the lung.
- **Intrapulmonary volume.** Intrapulmonary volume is the volume within the lungs. \bullet
- **Intrapleural pressure.** The normal pressure within the pleural space, the intrapleural pressure, is always negative, and this is the major factor preventing the collapse of the lungs.
- **Nonrespiratory air movements.** Nonrespiratory movements are a result of reflex activity, but some may be produced voluntarily such as <u>coughing</u>, sneezing, crying, laughing, hiccups, and yawning





Respiratory Volumes and Capacities







Respiratory Volumes and Capacities

Measurement	Adult male average value	Adult female average value	Description
Tidal volume (TV)	500 ml	500 ml	Amount of air inhaled or conditions
Inspiratory reserve volume (IRV)	3100 ml	1900 ml	Amount of air that can be volume inhalation
Expiratory reserve volume (ERV)	1200 ml	700 ml	Amount of air that can be volume exhalation
Residual volume (RV)	1200 ml	1100 ml	Amount of air remaining
Total lung capacity (TLC)6000 ml		4200 ml	Maximum amount of air of inspiratory effort: TLC =
Vital capacity (VC)	4800 ml	3100 ml	Maximum amount of air to inspiratory effort: VC = T
Inspiratory capacity (IC)	3600 ml	2400 ml	Maximum amount of air t expiration: IC = TV + IRV
Functional residual capacity (FRC)	2400 ml	1800 ml	Volume of air remaining expiration: FRC = ERV +



exhaled with each breath under resting

e forcefully inhaled after a normal tidal

e forcefully exhaled after a normal tidal

in the lungs after a forced exhalation

contained in lungs after a maximum TV + IRV + ERV + RV

that can be expired after a maximum V + IRV + ERV (should be 80% TLC)

that can be inspired after a normal

in the lungs after a normal tidal volume RV



Respiratory Sounds

- Bronchial sounds. Bronchial sounds are produced by air rushing through the large respiratory ulletpassageways (trachea and bronchi).
- Vesicular breathing sounds. Vesicular breathing sounds occur as air fills the alveoli, and they are soft • and resemble a muffled breeze.







External Respiration, Gas Transport and Internal Respiration

- External respiration. External respiration or pulmonary gas exchange involves oxygen being loaded and carbon dioxide being unloaded from the blood.
- **Internal respiration.** In internal respiration or systemic capillary gas exchange, oxygen is unloaded and carbon dioxide is loaded into the blood.
- **Gas transport.** Oxygen is transported in the blood in two ways: most attaches to hemoglobin molecules inside the RBCs to form oxyhemoglobin, or a very small amount of oxygen is carried dissolved in the plasma; while carbon dioxide is transported in plasma as bicarbonate ion, or a smaller amount (between 20 to 30 percent of the transported carbon dioxide) is carried inside the RBCs bound to hemoglobin





Control of Respiration

Neural Regulation

- **Phrenic and intercostal nerves.** These two nerves regulate the activity of the respiratory muscles, the diaphragm, and external intercostals.
- **Medulla and pons.** Neural centers that control respiratory rhythm and depth are located mainly in the ulletmedulla and pons; the medulla, which sets the basic rhythm of breathing, contains a pacemaker, or selfexciting inspiratory center, and an expiratory center that inhibits the pacemaker in a rhythmic way; pons centers appear to smooth out the basic rhythm of inspiration and expiration set by the medulla.
- **Eupnea.** The normal respiratory rate is referred to as eupnea, and it is maintained at a rate of **12 to 15** respirations/minute.
- **Hyperpnea.** During exercise, we breathe more vigorously and deeply because the brain centers send lacksquaremore impulses to the respiratory muscles, and this respiratory pattern is called hyperpnea.





Control of Respiration

Non-neural Factors Influencing Respiratory Rate and Depth

- **Physical factors.** Although the medulla's respiratory centers set the basic rhythm of breathing, there is no question that physical factors such as talking, coughing, and exercising can modify both the rate and depth of breathing, as well as an increased body temperature, which increases the rate of breathing.
- Volition (conscious control). Voluntary control of breathing is limited, and the respiratory centers will simply ignore messages from the cortex (our wishes) when the oxygen supply in the blood is getting low or blood pH is <u>falling</u>.
- **Emotional factors.** Emotional factors also modify the rate and depth of breathing through reflexes initiated by emotional stimuli acting through centers in the <u>hypothalamus</u>.
- Chemical factors. The most important factors that modify respiratory rate and depth are chemical- the levels of carbon dioxide and oxygen in the blood; increased levels of carbon dioxide and decreased blood pH are the most important stimuli leading to an increase in the rate and depth of breathing, while a decrease in oxygen levels become important stimuli when the levels are dangerously low.
- **Hyperventilation.** Hyperventilation blows off more carbon dioxide and decreases the amount of carbonic acid, • which returns blood pH to the normal range when carbon dioxide or other sources of acids begin to accumulate in the blood.
- Hypoventilation. Hypoventilation or extremely slow or shallow breathing allows carbon dioxide to accumulate in the blood and brings blood pH back into normal range when blood starts to become slightly alkaline.





Age-Related Physiological Changes in the Respiratory System

• Respiratory efficiency is reduced with age. They are unable to compensate for increased oxygen need and are significantly increasing the amount of air inspired. Therefore, difficulty in breathing is usually common especially during activities. Expiratory muscles become weaker so their <u>cough</u> efficiency is reduced and the amount of air left in the lungs is increased. <u>Health promotion</u> teaching can include smoking cessation, preventing respiratory infections through <u>handwashing</u>, and ensuring up to date <u>influenza</u> and <u>pneumonia</u> vaccinations.





THANK YOU

