



Topic: 1.11 – NATURE OF QUADRATIC FORM

Reduction of quadratic form to canonical form by orthogonal transformation - Nature of quadratic form.

Quadratic form:

A homogeneous polynomial of the second degree in any number of variables n called a quadratic form.

Eg: $2x_1^2 + 3x_2^2 - x_3^2 + 4x_1x_2 + 5x_1x_3 - 6x_2x_3$ is a quadratic form of three variables.

Note: The matrix corresponding to the quadratic form is

$$\begin{bmatrix} \text{coeff } x_1^2 & \frac{1}{2} \text{coeff. } x_1x_2 & \frac{1}{2} \text{coeff. } x_1x_3 \\ \frac{1}{2} \text{coeff. } x_2x_1 & \text{coeff. } x_2^2 & \frac{1}{2} \text{coeff. } x_2x_3 \\ \frac{1}{2} \text{coeff. } x_3x_1 & \frac{1}{2} \text{coeff. } x_3x_2 & \text{coeff. } x_3^2 \end{bmatrix}$$

Problem:

Write the matrix of the quadratic form
 $2x_1^2 - 2x_2^2 + 4x_3^2 + 2x_1x_2 - 6x_1x_3 + 6x_2x_3$



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

Solu:

$$Q = \begin{bmatrix} \text{coeff. } x_1^2 & \frac{1}{2} \text{coeff. } x_1 x_2 & \frac{1}{2} \text{coeff. } x_1 x_3 \\ \frac{1}{2} \text{coeff. } x_2 x_1 & \text{coeff. } x_2^2 & \frac{1}{2} \text{coeff. } x_2 x_3 \\ \frac{1}{2} \text{coeff. } x_3 x_1 & \frac{1}{2} \text{coeff. } x_3 x_2 & \text{coeff. } x_3^2 \end{bmatrix}$$

Here $x_2 x_1 = x_1 x_2$
 $x_3 x_1 = x_1 x_3$
 $x_2 x_3 = x_3 x_2$

$$\therefore Q = \begin{bmatrix} 2 & 1 & -3 \\ 1 & -2 & 3 \\ -3 & 3 & 4 \end{bmatrix}$$

write the matrix of the quadratic form
 $2x^2 + 8z^2 + 4xy + 10xz - 2yz$

Solu:

$$Q = \begin{bmatrix} \text{coeff. } x^2 & \frac{1}{2} \text{coeff. } x \cdot y & \frac{1}{2} \text{coeff. } x \cdot z \\ \frac{1}{2} \text{coeff. } y \cdot x & \text{coeff. } y^2 & \frac{1}{2} \text{coeff. } y \cdot z \\ \frac{1}{2} \text{coeff. } z \cdot x & \frac{1}{2} \text{coeff. } z \cdot y & \text{coeff. } z^2 \end{bmatrix}$$

$$Q = \begin{bmatrix} 2 & 2 & 5 \\ 2 & 0 & -1 \\ 5 & -1 & 8 \end{bmatrix}$$

write the quadratic form corresponding to the
following symmetric matrix $\begin{bmatrix} 0 & -1 & 2 \\ -1 & 1 & 4 \\ 2 & 4 & 3 \end{bmatrix}$

Solu:



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

∴ General form is

$$a_{11}x_1^2 + a_{22}x_2^2 + a_{33}x_3^2 + 2(a_{12})x_1x_2 + 2(a_{23})x_2x_3 + 2a_{13}x_1x_3 \\ = 5x_1^2 + x_2^2 + 3x_3^2 - 2x_1x_2 + 4x_1x_3 + 8x_2x_3$$

Quadratic form as a product of Matrices:

Let $A = [a_{ij}]$, $x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$ and $x^T = [x_1, x_2, \dots, x_n]$

∴ If we have a quadratic form $Q = \sum_{j=1}^n \sum_{i=1}^n a_{ij} x_i x_j$,
where $a_{ij} = a_{ji}$; then Q can be expressed as

$Q = x^T A x$ where the symmetric matrix
 $A = (a_{ij}) = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$ is called the
matrix of the Q.F.

Canonical form of a quadratic form.

Let $Q = x^T A x$ be a quadratic form in

n variables x_1, x_2, \dots, x_n - ^{original}
let $x = N y$ be a linear transformation,

where N is a normalized matrix

$$\text{Now } Q = x^T A x = (N y)^T A (N y) \\ = y^T (N^T A N) y$$



$$\begin{aligned} &= Y^T D Y \\ &= (y_1, y_2, \dots, y_n) \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} \\ &= \lambda_1 y_1^2 + \lambda_2 y_2^2 + \dots + \lambda_n y_n^2 \end{aligned}$$

(10) \therefore which is the canonical form of the quadratic form.

Note: This form is also called diagonalization of the quadratic form (10) to express the quadratic form as 'Sum of Squares'.

Nature of Quadratic forms:

Rank of A:
When the quadratic form is reduced to the canonical form it contains only r terms which is the rank of A .

Index of the Q.F (S):
The number of positive square terms in the canonical form is called the index of the quadratic form.

Signature of the Q.F
The difference of number of positive and negative square terms is called the signature of the quadratic form $[= S - (r - S) = 2S - r]$



Note :

The quadratic form $Q = x^T A x$ in n variables is said to be

(i) positive definite, if $r=n$ and $s=0$ (or) if all the eigenvalues of A are positive.

(ii) negative definite, if $r=0$ & $s=n$ (or) if all the eigenvalues of A are $-ve$.

(iii) positive Semidefinite, if $r \leq n$ and $s=0$
(or) if all the Eigenvalues of $A \geq 0$ & at least one Eigenvalue is zero.

(iv) Negative Semidefinite if $r=0$ & $s \leq n$ (or) if all the eigenvalues of $A \leq 0$ & at least one Eigenvalue is zero.

(v) Indefinite, in all other cases (or) if A has both $+ve$ & $-ve$ Eigenvalues.

Test for Nature of a Quadratic form through principal minors.

Let $A = [a_{ij}]$ be the matrix of the quadratic form in n variables x_1, x_2, \dots, x_n . Then A is a square symmetric matrix of order n .



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

Let $D_1 = |a_{11}|$
 $D_2 = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$
 $D_3 = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$
.....
 $D_n = |A|$

Here D_1, D_2, \dots, D_n are the principal minors of A .

(i) The Q.F is positive definite if D_1, D_2, \dots, D_n are all +ve i.e) $D_i > 0 \forall n$.

(ii) The Q.F is -ve definite if D_1, D_3, \dots are all -ve and D_2, D_4, D_6, \dots are all +ve

i.e) $(-1)^n D_n > 0 \forall n$.

(iii) The Q.F is +ve semi-definite if $D_n \geq 0$
& at least one $D_i = 0$

(iv) The Q.F is -ve semi-definite if $(-1)^n D_n \geq 0$
& at least one $D_i = 0$.

(v) The Q.F is indefinite in all other cases.