




AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

Topic: 3.3 – CENTRE OF CURVATURE





AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

1. Find the centre of curvature at the point $(am^2, 2am)$ on the parabola $y^2 = 4ax$.

Soln: Given $x = am^2$, $y = 2am$.

$$\frac{dx}{dm} = 2am; \quad \frac{dy}{dm} = 2a$$
$$y_1 = \frac{dy}{dm} \cdot \frac{dm}{dx} = \frac{2a}{2am} = \frac{1}{m}$$
$$y_2 = \frac{d}{dm} \left(\frac{dy}{dx} \right) \frac{dm}{dx} = \frac{d}{dm} \left(\frac{1}{m} \right) \frac{dm}{dx}$$
$$= -\frac{1}{m^2} \cdot \frac{1}{2am} = -\frac{1}{2am^3}$$
$$\bar{x} = x - \frac{y_1}{y_2} (1 + y_1^2)$$
$$= am^2 - \frac{1}{m} \left(\frac{-2am^3}{1} \right)^2 \left(1 + \frac{1}{m^2} \right)$$
$$= am^2 + 2am^2 (m^2 + 1)$$
$$\frac{\quad}{m^2} = am^2 + am^2 + 2a$$
$$= 3am^2 + 2a$$
$$\bar{y} = y + \frac{(1 + y_1^2)}{y_2}$$
$$= 2am + \frac{\left(1 + \frac{1}{m^2} \right)}{\frac{1}{2am^3}}$$
$$= 2am + \frac{(m^2 + 1)}{m^2} \cdot \frac{2am^3}{(-1)} = 2am - 2am^3 - 2am$$
$$\bar{y} = -2am^3$$

\therefore The centre of curvature is $(3am^2 + 2a, -2am^3)$.



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai


2) prove that if the centre of the curvature of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at one end of the minor axis lies at the other end, then the eccentricity of the ellipse is $\frac{1}{\sqrt{2}}$.

soln:

The ellipse is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \rightarrow \text{①}$

BB' is the minor axis. B is $(0, b)$

B' is $(0, -b)$



Diff. ① w.r.t x , we get

$$\frac{2x}{a^2} + \frac{2y}{b^2} \frac{dy}{dx} = 0.$$

$$y_1 = \frac{dy}{dx} = -\frac{2x}{a^2} \cdot \frac{b^2}{2y} = -\frac{b^2 x}{a^2 y}$$

$$y_2 = \frac{d^2 y}{dx^2} = -\frac{b^2}{a^2} \left[\frac{y(1) - x \cdot \frac{dy}{dx}}{y^2} \right]$$

$$y_1(0, b) = 0$$

$$y_2(0, b) = -\frac{b^2}{a^2} \left[\frac{b}{b^2} \right] = -\frac{b}{a^2}$$

Let (\bar{x}, \bar{y}) be the centre of curvature at

$$(0, b) \quad \bar{x} = x - \frac{y_1}{y_2} (1 + y_1^2) \quad \text{or} \quad \bar{y} = y + \frac{(1 + y_1^2)}{y_2}$$

$$\bar{x}(0, b) = 0 - 0 = 0$$



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

$$\bar{y}_{(9,m)} = b + \frac{a^2}{-b^2} (1+0) = b - \frac{a^2}{b}$$

The centre of curvature is $(0, b - \frac{a^2}{b})$ this is given to be the point $(0, -b)$ the other end of the minor axis.

$$b - \frac{a^2}{b} = -b \Rightarrow b^2 - a^2 = -b^2$$

$$2b^2 = a^2 \rightarrow \textcircled{2}$$

But $b^2 = a^2(1-e^2)$ where e is being eccentricity using in $\textcircled{2}$

$$a^2 = 2a^2(1-e^2)$$

$$1-e^2 = \frac{1}{2} \Rightarrow e^2 = \frac{1}{2}$$

$$e = \frac{1}{\sqrt{2}}$$

centre of curvature:

The centre of curvature (\bar{x}, \bar{y}) at any point $P(\bar{x}, \bar{y})$ on the curve $y=f(x)$ are

$$\bar{x} = x - \frac{y_1}{\frac{y_2}{2}} (1+y_1^2)$$

$$\bar{y} = y + \frac{1}{\frac{y_2}{2}} (1+y_1^2)$$