



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

Topic: 1.6 – DIAGONALIZATION OF MATRICES

Working Rule For Diagonalisation:

- * To find the characteristic equation.
- * To solve the characteristic equation.
- * To find the Eigen values
- * To find the Eigen vectors
- * If the Eigen vectors are orthogonal, then form a normalized modal N .
- * Normalized form is $N = \begin{bmatrix} x_1/l \\ x_2/l \\ x_3/l \end{bmatrix}$ where $l = \sqrt{x_1^2 + x_2^2 + x_3^2}$
- * Find N^T
- * Calculate AN
- * Calculate $D = N^TAN$

Problems:

1. Diagonalise the matrix $\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$

Let $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$

Characteristic equation $\Rightarrow \lambda^3 - 18\lambda^2 + 45\lambda = 0$

Eigen values are 0, 3, 15

- Eigen vectors
- i) If $\lambda = 0$, then Eigen vector is $[1 \ 2 \ 2]^T$
 - ii) If $\lambda = 3$, then Eigen vector is $[2 \ 1 \ -2]^T$
 - iii) If $\lambda = 15$, then Eigen vector is $[2 \ -2 \ 1]^T$



AN AUTONOMOUS INSTITUTION

Accredited by NBA – AICTE and Accredited by NAAC – UGC with 'A' Grade
Approved by AICTE, New Delhi & Affiliated to Anna University, Chennai

To find the Eigen vectors are orthogonal to each other.

$$X_1^T X_2 \Rightarrow 2+2-4 = 0$$

$$X_2^T X_3 \Rightarrow 4-2-2 = 0$$

$$X_1^T X_3 \Rightarrow 2-4+2 = 0$$

They are orthogonal to each other.

To form the Normalised Matrix:

$$N = \begin{bmatrix} 1/3 & 2/3 & 2/3 \\ 2/3 & 1/3 & -2/3 \\ 2/3 & -2/3 & 1/3 \end{bmatrix} \Rightarrow \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$$

To find the transpose of normalised matrix

$$N^T = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \frac{1}{3}$$

Calculate AN

$$AN = \frac{1}{3} \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 10 \\ 0 & 1 & -10 \\ 0 & -2 & 5 \end{bmatrix}$$

Calculate Diagonalised Matrix D

$$D = NAN^T$$

$$= \frac{1}{3} \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \cdot \frac{1}{3}$$

$$= \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 & 10 \\ 0 & 1 & -10 \\ 0 & -2 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 15 \end{bmatrix}$$