



SNS COLLEGE OF NURSING

saravanampatti(po),coimbatore-35 Affiliated to The Tamilnadu DR.MGR MEDICAL UNIVERSITY, Chennai DEPARTMENT OF NURSING COURSE – 1 ST YEAR B.SC(N) **Subject-Nursing Foundation UNIT: X- MEETING NEEDS OF PATIENT TOPIC 2: Providing urinal/Bed pan/condom drinage**

























Meeting elimination need of bedridden male patients using a urinal.

Meeting elimination need of bedridden female patients using a bedpan.









- \checkmark To facilitate bowel and bladder elimination
- ✓ To collect specimen
- \checkmark To give perineal wash
- ✓ To perform bowel and bladder training







- Urinal or bedpan
- Paper or washable cover
- Toilet tissue
- Towel Plastic or rubber bed protector Gloves



PROCEDURE-URINAL



PROCEDURE	RATIONAL
Assess patients normal	Identifies normal pattern
elimination habits	of urination
Palpate for distended bladder	Identifies if bladder is full
	and patient needs to void
Wash hands and wear gloves	Reduces transmission of
	micro-organisms





•Provide privacy by closing the door or bedside screens	Provides comfort
Ask the patient to sit.	Easier to void
Patient should hold the	avoids spillage of urine on
urinal and position penis in	the bed linen
the urinal and to void.	





	 _	_
Drovida	10 1h0	notiont
Provide		

Once patient has finished	
remove the urinal	
If required collect urine in the specimen container	For testing of urine
Empty the urinal into toilet and	
clean it.	



PROCEDURE-URINALS



Empty the urinal into	
toilet and clean it	
 Place urinal near to 	
patient after cleaning	
for next use	
Remove gloves and	
wash hands.	



PROVIDING URINAL





PROVIDIND URINAL/BED PAN/CONDOM DRINAGE





PROCEDURE	RATIONALE
Assess the patients level of mobility, amount of assistance required .	•Helps in providing assistance required by the nurse
Wash hands and wear gloves	 Reduces transmission of micro-organisms



PROCEDURE-BED PAN



Warm bedpan under warm	•Warm pan helps to
water for few seconds	relax anal sphincter
and dry it	
 Provide privacy by closing the door or bedside screens 	 Provides comfort
the door of bedside screens	
Position bed to convenient	 Prevents strain on
height	back



PROCEDURE-BED PAN



•Elevate the side rails

•Reduces the risk of

accidental falls

Ask the patient to raise the hips and place the bed pan.

 If the patient is unconscious, position patient to side-lying position, place bedpan against buttocks and roll the patient to the bed pan.





If patient can help by himself, provide tissue to wipe or place the another bed pan and pour water to clean himself.

 Provide soap and water to clean patients hand. If patient is helpless, provides perineal care and anal care

•Send the bedpan to clean

Remove gloves and wash hands



PROVIDING -BED PAN







AFTER CARE



Document in patient's record:

- Time and results of elimination.
- Assistance required.
- Report any changes in the patient's condition to supervisor.



CONDOM DRINAGE







STEP1:MEASUREMENT



• If you are unsure of the size of

condom to select, use a

measuring guide.

Place the guide behind the

glands of the penis. If between sizes, select a larger size.









- Wash your hands and your penis with soap and water.
- Rinse and dry your penis thoroughly. Before applying the sheath, trim your pubic hair.
- Alternatively, use a paper towel or piece of fabric with a hole in the middle as a guard to keep the pubic hair away.



STEP 2 PREPARATION



 Inspect penis to make sure there's no broken or reddened skin and don't use lotions, oils as they may prevent a secure seal against the skin.









- If not circumcised make sure the foreskin is pulled forward over the head of the penis before applying the sheath.
- Don't push the sheath hard against the penis tip.
 Leave a gap of approx 5cm. Once rolled down, use the hand to make sure the condom is fitted correctly.





- If you're in bed, you can attach the collection bag to your bed. If you're sitting in a chair, you can attach the urine bag to the chair below the cushion or seat.
- Connect the condom to the tube of the urine bag.





- If you're using a small leg bag, use a catheter leg strap to secure the urine bag to your leg just below your knee.
- Leave some slack in the tube so the condom will not be pulled when you move your leg .The urine collection bag must be placed so that your urine flows downward.



APPLICATION (continued)









- Remove the sheath whilst showering. (Adding a few drops of shampoo/conditioner make removal easier).
- Ensure to clean under the foreskin. If there is a problem with the sheath coming off when using an overnight drainage bag, check the bag tubing width.
 Try a bag with a wider tube.





- 1.What is bedpan and urinal?
- 2.What all are the purpose of giving bedpan?
- 3.What are all the articles required for this procedure?
- 4. How to apply condom catheter?



REFERENCES



- Helen Harkreader. Fundamentals of nursing caring and clinical judgement. Elsevier publication St. Louis: 2009;(3).
- Annamma Jacob, Rekha R. clinical nursing procedures the art of nursing practice. Jaypee publications Newdelhi: 2015