



DEPARTMENT OF MATHEMATICS

UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

PARTIAL DERIVATIVES :

Let $u = f(x, y)$ be a function of two independent variables. Differentiating 'u' w.r. to 'x' keeping 'y' as constant is known as partial derivative of u w.r. to x and is denoted by $\frac{\partial u}{\partial x}$ (i) u_x .

Similarly, $\frac{\partial u}{\partial y}$ (ii) u_y .

NOTE :

(i) If $u = v + w + \dots$ where v, w are all functions of x, y, z, \dots then.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial x} + \frac{\partial w}{\partial x} + \dots$$

$$\frac{\partial u}{\partial y} = \frac{\partial v}{\partial y} + \frac{\partial w}{\partial y} + \dots$$

(ii) If u & v are functions of x, y, z etc. then

$$\frac{\partial}{\partial x} (uv) = u \frac{\partial v}{\partial x} + v \frac{\partial u}{\partial x}$$

$$\frac{\partial}{\partial y} (uv) = u \frac{\partial v}{\partial y} + v \frac{\partial u}{\partial y}$$

(iii) If u & v are functions of x, y, z etc. then

$$\frac{\partial}{\partial x} \left(\frac{u}{v} \right) = \frac{v \frac{\partial u}{\partial x} - u \frac{\partial v}{\partial x}}{v^2}$$

$$\frac{\partial}{\partial y} \left(\frac{u}{v} \right) = \frac{v \frac{\partial u}{\partial y} - u \frac{\partial v}{\partial y}}{v^2}$$



DEPARTMENT OF MATHEMATICS

UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

(iv) If u is a function of t where t is a function of the variables x, y, z, \dots then.

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial t} \cdot \frac{\partial t}{\partial x}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial t} \cdot \frac{\partial t}{\partial y} \dots \dots$$

SUCCESSIVE PARTIAL DIFFERENTIATION:

Let $z = f(x, y)$ then $\frac{\partial z}{\partial x}$ & $\frac{\partial z}{\partial y}$ being the function of x & y can be further be differentiated partially w.r.t. to x & y .

\therefore we have $\frac{\partial^2 z}{\partial x^2}$, $\frac{\partial^2 z}{\partial y^2}$, $\frac{\partial^2 z}{\partial x \partial y}$, $\frac{\partial^2 z}{\partial y \partial x}$

Note: $\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$.

① If $u = \frac{x}{y} + \frac{y}{z} + \frac{z}{x}$, find $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z}$.

Sol: $u = \frac{x}{y} + \frac{y}{z} + \frac{z}{x}$.



DEPARTMENT OF MATHEMATICS

UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

Soln: $u = \frac{x}{y} + \frac{y}{z} + \frac{z}{x}$

$$\frac{\partial u}{\partial x} = \frac{1}{y} - \frac{z}{x^2} \Rightarrow x \frac{\partial u}{\partial x} = \frac{x}{y} - \frac{z}{x}$$

$$\frac{\partial u}{\partial y} = -\frac{x}{y^2} + \frac{1}{z} \Rightarrow y \frac{\partial u}{\partial y} = -\frac{x}{y} + \frac{y}{z}$$

$$\frac{\partial u}{\partial z} = -\frac{y}{z^2} + \frac{1}{x} \Rightarrow z \frac{\partial u}{\partial z} = -\frac{y}{z} + \frac{z}{x}$$

$$\therefore x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$$

② If $u = (x-y)^2 + (y-z)^2 + (z-x)^2$. p.t. $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$

Soln: $u = (x-y)^2 + (y-z)^2 + (z-x)^2$

$$\frac{\partial u}{\partial x} = 2(x-y)(1) + 2(z-x)(-1) = 2(x-y) - 2(z-x)$$

$$\frac{\partial u}{\partial y} = 2(y-z)(1) + 2(x-y)(-1) = 2(y-z) - 2(x-y)$$

$$\frac{\partial u}{\partial z} = 2(z-x)(1) + 2(y-z)(-1) = 2(z-x) - 2(y-z)$$

$$\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$$



DEPARTMENT OF MATHEMATICS

UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

$$Q) \text{ If } r^2 = (x-a)^2 + (y-b)^2 + (z-c)^2 \text{ s.t. } \frac{\partial^2 r}{\partial x^2} + \frac{\partial^2 r}{\partial y^2} + \frac{\partial^2 r}{\partial z^2} = \frac{2}{r}$$

Soln: $r^2 = (x-a)^2 + (y-b)^2 + (z-c)^2$

$$2r \cdot \frac{\partial r}{\partial x} = 2(x-a)$$

d.w.r to x $\frac{\partial r}{\partial x} = \frac{(x-a)}{r}$

$$\frac{\partial^2 r}{\partial x^2} = \frac{r \cdot 1 - (x-a) \frac{\partial r}{\partial x}}{r^2} = \frac{r - (x-a) \cdot \frac{(x-a)}{r}}{r^2}$$

$$= \frac{r^2 - (x-a)^2}{r^3}$$

p.d.w.r to y $\frac{\partial r}{\partial y} = \frac{y-b}{r}$

$$\frac{\partial^2 r}{\partial y^2} = \frac{r^2 - (y-b)^2}{r^3}$$

p.d.w.r to z $\frac{\partial r}{\partial z} = \frac{z-c}{r}$

$$\frac{\partial^2 r}{\partial z^2} = \frac{r^2 - (z-c)^2}{r^3}$$

$$\frac{\partial^2 r}{\partial x^2} + \frac{\partial^2 r}{\partial y^2} + \frac{\partial^2 r}{\partial z^2} = \frac{r^2 - (x-a)^2 + r^2 - (y-b)^2 + r^2 - (z-c)^2}{r^3}$$

$$= \frac{3r^2 - [(x-a)^2 + (y-b)^2 + (z-c)^2]}{r^3} = \frac{3r^2 - r^2}{r^3}$$

$$= \frac{2}{r}$$