



DEPARTMENT OF MATHEMATICS UNIT – II DESIGN OF EXPERIMENTS

UNIT-II DESIGN OF EXPERIMENTS

ANALYSIS OF VARIANCE (ANOVA):

ANOVA is a technique that will enable us to test the significance of the difference among more than two sample mean.

ASSUMPTION:

- 1) The observations are random.
- 2) The observations are independent.
- 3) The samples are drawn from normal populations.
- 4) Population variances are equal.

BASIC PRINCIPLES:

- 1) Randomisation
- 2) Replication
- 3) Local control.

BASIC DESIGN:

- * Completely randomised design (CRD) One-way classification
- * Randomised Block design (RBD) Two-way classification
- * Latin square design (LSD) Three-way classification
- * Two square factorial design

Hint :- F-Ratio : $F = \frac{S_1^2}{S_2^2}$ where $S_1^2 > S_2^2$



DEPARTMENT OF MATHEMATICS

UNIT - II DESIGN OF EXPERIMENTS

procedure to find :-

- 2) Sum of all the terms (T) & total no of sample size (N)
- 3) Correction factor ($C.F$), $C.F = \frac{T^2}{N}$
- 4) TSS : Total sum of squares
= (sum of the squares of all the terms) - C.F.
- 5) SSC : Sum of squares between samples
- 6) SSE : Error sum of squares
= TSS - SSC
- 7) Anova table

8) Conclusion :

1) $H_0: \mu_1 = \mu_2 = \dots = \mu_k$

1) A completely randomised design experiment with 10 plots and 3 treatments gave the following result :

plot No. :	1	2	3	4	5	6	7	8	9	10
treatment :	A	B	C	A	C	C	A	B	A	B
yield :	5	4	3	7	5	1	3	4	1	4

Analyse the result for treatment effects.



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore - 35



DEPARTMENT OF MATHEMATICS

UNIT - II DESIGN OF EXPERIMENTS

Treatment	Yield				Treatment
	A	B	C		A B C
(n ₁) A	5	4	3	1	5 4 3
(n ₂) B	4	4	4	-	4 4 5
(n ₃) C	3	5	1	-	3 4 1

x ₁	x ₂	x ₃	Total	x ₁ ²	x ₂ ²	x ₃ ²
5	4	3	12	25	16	9
4	4	5	16	16	16	25
3	4	1	11	9	16	1
1	-	-	1	1	-	-
$\frac{16}{\Sigma n_1}$	$\frac{15}{\Sigma n_2}$	$\frac{9}{\Sigma n_3}$	40	$\frac{84}{\Sigma n_1^2}$	$\frac{81}{\Sigma n_2^2}$	$\frac{35}{\Sigma n_3^2}$

Step 1: Formulating H₀ & H₁:

H₀: there is no significance difference between the treatments.

H₁: there is significance difference between the treatments.

Step 2: To find T & N:

$$T = \Sigma n_1 + \Sigma n_2 + \Sigma n_3$$

$$= 16 + 15 + 9 = 40$$

$$N = n_1 + n_2 + n_3$$

$$= 4 + 3 + 3 = 10$$



DEPARTMENT OF MATHEMATICS

UNIT - II DESIGN OF EXPERIMENTS

Step 3: Correction Factor, C.F.

$$C.F = \frac{T^2}{N} = \frac{40^2}{10} = 160$$

$$\begin{aligned} \text{Step 4: } TSS &= \sum n_1^2 + \sum n_2^2 + \sum n_3^2 - C.F \\ &= 84 + 81 + 35 - 160 \\ &= 40 \end{aligned}$$

$$\begin{aligned} \text{Step 5: } SSC &= \frac{(\sum x_1)^2}{n_1} + \frac{(\sum x_2)^2}{n_2} + \frac{(\sum x_3)^2}{n_3} - C.F. \\ &= \frac{16^2}{4} + \frac{15^2}{3} + \frac{9^2}{3} - 160 \\ &= 6 \end{aligned}$$

$$\begin{aligned} \text{Step 6: } SSE &= TSS - SSC \\ &= 40 - 6 = 34 \end{aligned}$$

Step 7: Anova table:

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	F-Rat
Between samples (Column)	SSC : 6	C-1 = 3-1 = 2	MSC = $\frac{6}{2} = 3$	$F_c = \frac{4}{3} = 1.33$
With samples (Error)	SSE : 34	N-c = 10-3 = 7	MSE = $\frac{34}{7} = 4.86$	$F_{0.05} = 7.7$

Step 8: Conclusion:

$$F_c = 1.33 < 7.7 = F_{\alpha}, H_0 \text{ is accepted.}$$

(a) There is no significant difference between the treatments.