



DEPARTMENT OF MATHEMATICS


(32)

TYPE III : Problems on Steady State conditions and non-zero boundary conditions (or) Non-zero temperatures at the ends of the bar, both in steady state and unsteady state :

- ① Two ends A and B of a rod of length 20 cms have their temperatures at 30°C and 80°C respectively until steady state conditions prevail. Then the temperatures at the ends A and B are changed to 40°C and 60°C respectively. Find the temperature distribution of the rod at any time 't'.

Solution :

Steady State 1 :



The steady state solution is

$$u = ax + b$$

(i) $u(0) = 30^\circ\text{C}$

(ii) $u(20) = 80^\circ\text{C}$

$$u(0) = a(0) + b = 30$$

$$\boxed{b = 30}$$

$$u(20) = 20a + b = 80$$

$$20a + 30 = 80$$

$$20a = 50$$

$$\boxed{a = 5/2}$$

$$\therefore \boxed{u(x) = \frac{5}{2}x + 30}$$

which is the initial temperature of the rod. i.e.,

$$u(x, 0) = \frac{5}{2}x + 30$$




SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)
DEPARTMENT OF MATHEMATICS



Steady state 2 :



The solution is, $u(x) = ax + b \rightarrow \textcircled{1}$

(i) $u(0) = 40^\circ\text{C}$

(ii) $u(20) = 60^\circ\text{C}$

App (i) $\Rightarrow u(0) = a(0) + b = 40$
in $\textcircled{1}$ $b = 40$

App (ii) in $\textcircled{1}$ $\Rightarrow u(20) = 20a + b = 60$
 $20a + 40 = 60$
 $a = 1$

$\therefore u(x) = x + 40$

which is the transient state temperature of the rod.

i.e., $u_t(x, t) = x + 40$.

Step 1 : The one-dimensional heat equation is,

$$\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$$

Step 2 : The boundary conditions are,

(i) $u(0, t) = 40$

(ii) $u(20, t) = 60$

(iii) $u(x, 0) = \frac{5}{2}x + 30$

Step 3 : The correct solution is,

$$u(x, t) = (c_1 \cos px + c_2 \sin px) e^{-\alpha^2 p^2 t} + u_t$$

$$u(x, t) = x + 40 + (c_1 \cos px + c_2 \sin px) e^{-\alpha^2 p^2 t} \rightarrow \textcircled{1}$$

Step 4 : Applying Condition (i) in $\textcircled{1}$,

$$u(0, t) = 40 + c_1 e^{-\alpha^2 p^2 t}$$

$$40 = 40 + c_1 e^{-\alpha^2 p^2 t}$$

$$0 = c_1 e^{-\alpha^2 p^2 t}$$



Here $e^{-\alpha^2 p^2 t} \neq 0$ [\because It is defined for all t]

$$\therefore C_1 = 0$$

$$\textcircled{1} \Rightarrow u(x, t) = x + 40 + C_2 \sin px e^{-\alpha^2 p^2 t} \rightarrow \textcircled{2}$$

Step 5: Applying condition (ii) in $\textcircled{2}$.

$$u(20, t) = 20 + 40 + C_2 \sin 20p e^{-\alpha^2 p^2 t}$$

$$60 = 60 + C_2 \sin 20p e^{-\alpha^2 p^2 t}$$

$$0 = C_2 \sin 20p e^{-\alpha^2 p^2 t}$$

Here $e^{-\alpha^2 p^2 t} \neq 0$ [\because It is defined for all t]

$C_2 \neq 0$ [\because $C_1 = 0$ we get a trivial solution]

$$\therefore \sin 20p = 0$$

$$\sin 20p = \sin n\pi$$

$$20p = n\pi$$

$$p = \frac{n\pi}{20}$$

$$\textcircled{2} \Rightarrow u(x, t) = x + 40 + C_2 \sin\left(\frac{n\pi x}{20}\right) e^{-\alpha^2 n^2 \pi^2 t / 20^2}$$

$$u(x, t) = x + 40 + C_n \sin\left(\frac{n\pi x}{20}\right) e^{-\alpha^2 n^2 \pi^2 t / 400}$$

Step 6: The most general solution is,

$$u(x, t) = x + 40 + \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{20}\right) e^{-\alpha^2 n^2 \pi^2 t / 400} \rightarrow \textcircled{3}$$

Step 7: Applying condition (iii) in $\textcircled{3}$,

$$u(x, 0) = x + 40 + \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{20}\right)$$

$$\frac{5}{2}x + 30 = x + 40 + \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{20}\right)$$

$$\frac{5}{2}x - x + 30 - 40 = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{20}\right)$$

$$\frac{3}{2}x - 10 = \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi x}{20}\right) \rightarrow \textcircled{4}$$