Reduction of quadratic form to canonical form by orthogonal transformation

Reduce the quadratic form $2x_1^2 + 2x_2^2 + x_3^2 + 4x_1x_2 = 0$ to canonical form by orthogonal reduction .Find rank, index, signature and nature

<u>Step 1:</u>

The matrix form is

$$A = \begin{bmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

<u>Step 2:</u>

Characteristic equation ,Eigen values, Eigen vectors

C1 =Sum of leading diagonal elements

=2+2+1 =5

 C_2 = Sum of minors of leading diagonal elements

=4

 $C_3 = |A|$

 $=\begin{vmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{vmatrix}$ = 0

The characteristic equation is

$$\lambda^3 - 5\lambda^2 + 4\lambda = 0$$

The eigen values are 0,1,4

The eigen vectors are $(A - \lambda I)X = 0$

Reduction of quadratic form to canonical form by orthogonal transformation

$$\begin{pmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \Big] \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$
$$\begin{pmatrix} 2 -\lambda & 2 & 0 \\ 2 & 2 -\lambda & 0 \\ 0 & 0 & 1 - \lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{pmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

CASE (i)

When $\lambda = 0$

$$\begin{pmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

The cofactor of first row elements are $\begin{pmatrix} 2 \\ -2 \\ 0 \end{pmatrix}$ ie $\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$

The Eigen vector when $\lambda = 0$ is $\begin{pmatrix} -1 \\ 0 \end{pmatrix}$

CASE (ii)

When $\lambda = 1$

$$\begin{pmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

The cofactor of third row elements are $\begin{pmatrix} 0 \\ 0 \\ -3 \end{pmatrix}$ ie $\begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$

Reduction of quadratic form to canonical form by orthogonal transformation

The Eigen vector when
$$\lambda = 1$$
 is $\begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$

CASE (iii)

When $\lambda = 4$

$$\begin{pmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

The cofactor of first row elements are $\begin{pmatrix} 6 \\ 6 \\ 0 \end{pmatrix}$ ie $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$
The Eigen vector when $\lambda = 4$ is $\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$

STEP 3:

To check pair wise orthogonality

$$X_{1}^{T}X_{2} = (2 -2 0) \begin{pmatrix} 0\\0\\-1 \end{pmatrix} = 0$$
$$X_{2}^{T}X_{3} = (0 0 -1) \begin{pmatrix} 1\\1\\0 \end{pmatrix} = 0$$
$$X_{3}^{T}X_{1} = (1 1 0) \begin{pmatrix} 2\\-2\\0 \end{pmatrix} = 0$$

STEP 4:

To find normalized vector

Reduction of quadratic form to canonical form by orthogonal transformation

Eigen vector	$l(x) = \sqrt{x_1^2 + x_2^2 + x_3^2}$	Normalized vector = $\begin{pmatrix} x_1/l(x_1) \\ x_2/l(x_2) \\ x_3/l(x_3) \end{pmatrix}$
$\begin{pmatrix} 1\\ -1\\ 0 \end{pmatrix}$	$\sqrt{1+1+0} = \sqrt{2}$	$\begin{pmatrix} 1//\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix}$
$\begin{pmatrix} 0\\ 0\\ -1 \end{pmatrix}$	$\sqrt{0+0+1} = \sqrt{1}$	$\begin{pmatrix} 0\\ 0\\ -1 \end{pmatrix}$
$\begin{pmatrix} 1\\1\\0 \end{pmatrix}$	$\sqrt{1+1+0} = \sqrt{2}$	$\begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{pmatrix}$

<u>STEP 5:</u>

Normalized modal matrix

$$\mathbf{N} = \begin{bmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & -1 & 0 \end{bmatrix}$$

$$\mathbf{N}^{\mathrm{T}} = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0\\ 0 & 0 & -1\\ 1/\sqrt{2} & 1/\sqrt{2} & 1 \end{bmatrix}$$

<u>STEP 6:</u>

$$NN^T = N^T N = I$$

Reduction of quadratic form to canonical form by orthogonal transformation

$$N^{T}N = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0\\ 0 & 0 & -1\\ 1/\sqrt{2} & 1/\sqrt{2} & 1 \end{pmatrix} \begin{pmatrix} 1//\sqrt{2} & 0 & 1/\sqrt{2}\\ -1/\sqrt{2} & 0 & 1/\sqrt{2}\\ 0 & -1 & 0 \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{pmatrix}$$
$$= I$$

STEP 7:

To find diagonalyze matrix

$$N^{T}AN = D$$

$$N^{T}A = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0\\ 0 & 0 & -1\\ 1/\sqrt{2} & 1/\sqrt{2} & 1 \end{pmatrix} \begin{pmatrix} 2 & 2 & 0\\ 2 & 2 & 0\\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0\\ 0 & 0 & -1\\ 4/\sqrt{2} & 4/\sqrt{2} & 0 \end{pmatrix}$$

$$N^{T}AN = \begin{pmatrix} 0 & 0 & 0\\ 0 & 0 & -1\\ 4/\sqrt{2} & 4/\sqrt{2} & 0 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2}\\ -1/\sqrt{2} & 0 & 1/\sqrt{2}\\ 0 & -1 & 0 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 4 \end{pmatrix}$$

Reduction of quadratic form to canonical form by orthogonal transformation

Step 8:

 $Y^T D Y = 0$

 $0y_1^2 + y_2^2 + 4y_3^2 = 0$

The index p=2

Rank r=2

Signature s=2p-r =2

The nature is semi positive