



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)
Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

Circle of Curvature:

The circle of curvature of a curve $y = f(x)$ at a point $P(x, y)$ with the centre of curvature, $C(\bar{x}, \bar{y})$ and radius curvature ρ is

$$(x - \bar{x})^2 + (y - \bar{y})^2 = \rho^2$$

1. Find the circle of curvature of the curve $x^3 + y^3 = 3axy$ at the

$$point \left(\frac{3a}{2}, \frac{3a}{2} \right)$$

Soln :

$$\text{Given } x^3 + y^3 = 3axy \quad \dots (1)$$

Differentiating (1) w.r.to. 'x', we get

$$3x^2 + 3y^2 \frac{dy}{dx} = 3a [x \frac{dy}{dx} + y]$$

$$\Rightarrow \frac{dy}{dx} = \frac{ay - x^2}{y^2 - ax} \quad \dots (2)$$

$$y_1 = \left(\frac{dy}{dx} \right)'_{\left(\frac{3a}{2}, \frac{3a}{2} \right)}$$

$$= \frac{\frac{3a}{2} - 9a^2}{\frac{9a^2}{4} - a \frac{3a}{2}} - \frac{\frac{3a^2}{2} - 9a^2}{\frac{9a^2}{4} - a \frac{3a^2}{2}}$$

$$= -1$$

$$y_1 = -1$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

$$From \ (2), \frac{d^2y}{dx^2} = \frac{(y^2 - ax) \left(a \frac{dy}{dx} - 2x \right) - (ay - x^2) \left(2y \frac{dy}{dx} - a \right)}{(y^2 - ax)^2}$$

$$\begin{aligned} y_2 &= \left(\frac{d^2y}{dx^2} \right)_{\substack{3a \\ (2, 2)}} \\ &= \frac{\left(\frac{9a^2}{4} - \frac{3a^2}{2} \right) \left(-a - \frac{6a}{2} \right) - \left(\frac{3a^2}{2} - \frac{9a^2}{4} \right) \left(-\frac{6a}{2} - a \right)}{\left(\frac{9a^2}{4} - \frac{3a^2}{2} \right)^2} \\ &= \frac{(-4a) - 4a}{\frac{9a^2}{4} - \frac{3a^2}{2}} \\ &= \frac{-8a}{\frac{3a^2}{4}} = \frac{-32}{3a} \end{aligned}$$

$$y_2 = \frac{-32}{3a}$$

\therefore Radius of curvature

$$\begin{aligned} \therefore \rho &= \frac{\left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}} = \frac{\left[1 + (-1)^2 \right]^{\frac{3}{2}}}{\frac{-32}{3a}} \\ &= \frac{(2)^{\frac{3}{2}} \times 3a}{-32} \\ &= \frac{3\sqrt{2} \times a}{16} \end{aligned}$$

\therefore The centre of curvature



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

$$x = x - y_1 \left(\frac{1+y_1^2}{y_2} \right)$$

$$= \frac{3a}{2} - \frac{(-1)(2)}{-32} 3a = \frac{21a}{16}$$

$$= \frac{3a}{2} + \frac{(2)}{-32} \cdot 3a = \frac{21a}{16}$$

\therefore The required circle of curvature is

$$(x - \frac{21a}{16})^2 + (y - \frac{21a}{16})^2 = (\frac{3\sqrt{2} \times a}{16})^2$$

$$\text{i.e., } (x^2 + y^2) - \frac{21a}{8}(x + y) + \frac{432a^2}{128} = 0$$

2. Find the circle of curvature of the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ **at the point** $\left(\frac{a}{4}, \frac{a}{4}\right)$.

Soln:

Given curve: $\sqrt{x} + \sqrt{y} = \sqrt{a}$ ----> (1)

Point: $P\left(\frac{a}{4}, \frac{a}{4}\right)$

Centre of curvature $C(\bar{x}, \bar{y})$,

$$x = x - y_1 \left(\frac{1+y_1^2}{y_2} \right)$$

$$y = y + \left(\frac{1+y_1^2}{y_2} \right)$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

From (1),

$$\sqrt{x} + \sqrt{y} = \sqrt{a} \Rightarrow \text{Differentiate w.r.to. } x$$

$$\frac{d}{dx}(\sqrt{x} + \sqrt{y}) = \frac{d}{dx}(\sqrt{a})$$

$$\frac{1}{2}x^{-\frac{1}{2}} + \frac{1}{2}y^{-\frac{1}{2}}\frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{x^{-\frac{1}{2}}}{y^{-\frac{1}{2}}} = -\frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}}$$

$$\text{at } P\left(\frac{a}{4}, \frac{a}{4}\right), \frac{dy}{dx} = -1$$

$$y_1 = -1$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx}\left(\frac{dy}{dx}\right) \\ &= \frac{d}{dx}\left(-\frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}}\right) \\ &= -\left[\frac{x^{\frac{1}{2}}\left(\frac{1}{2}y^{-\frac{1}{2}}\right)\left(\frac{dy}{dx}\right) - y^{-\frac{1}{2}}\left(\frac{1}{2}x^{-\frac{1}{2}}\right)}{x}\right] \\ &= -\frac{1}{2}\left[\frac{x^{\frac{1}{2}}y^{-\frac{1}{2}}\frac{dy}{dx} - y^{-\frac{1}{2}}x^{-\frac{1}{2}}}{x}\right] \end{aligned}$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

$$\text{At } P\left(\frac{a}{4}, \frac{a}{4}\right), \frac{d^2y}{dx^2} = \frac{-1}{2} \left[\frac{(1)(-1) - (1)}{\frac{a}{4}} \right]$$

$$= \frac{-1}{2} \left[\frac{-2}{\frac{a}{4}} \right]$$

$$y_2 = \frac{4}{a}$$

$$\frac{1+y_1^2}{y_2} = \frac{1+(-1)^2}{\frac{a}{4}} = \frac{2}{\frac{a}{4}} = \frac{a}{2}$$

Consider

$$x = \frac{a}{4} - (-1) \left(\frac{a}{2} \right) = \frac{a}{4} + \frac{a}{2} = \frac{3a}{4}$$

$$\bar{y} = \frac{a}{4} + \frac{a}{2} = \frac{3a}{4}$$

We know that

$$\rho = \frac{(1+y^2)^{3/2}}{y_2}$$

$$\therefore \text{At } P\left(\frac{a}{4}, \frac{a}{4}\right), \rho = \frac{\left[1+(-1)^2\right]^{3/2}}{\frac{a}{4}} = (2)^{3/2} \times \frac{a}{4}$$

$$= 2\sqrt{2} \times \frac{a}{4}$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

$$= \frac{a\sqrt{2}}{2}$$

$$\therefore \rho = \frac{a\sqrt{2}}{2}$$

From (2), we have

$$\left(x - \frac{3a}{4}\right)^2 + \left(y - \frac{3a}{4}\right)^2 = \left(\frac{a}{\sqrt{2}}\right)^2,$$

3. Find the equation of the circle of curvature at (c, c) on $xy = c^2$.

Soln:

$$\text{Given curve: } xy = c^2 \quad \longrightarrow \quad (1)$$

Point: $P(c, c)$

Centre of curvature $C(\bar{x}, \bar{y})$,

$$\begin{aligned} x &= x - y_1 \left(\frac{1+y_1^2}{y_2} \right) \\ \bar{y} &= y + \left(\frac{1+y^2}{y_2} \right) \quad \text{dy} \quad d^2y \\ &\text{where } y_1 = \frac{dy}{dx}; \quad y_2 = \frac{d^2y}{dx^2}. \end{aligned}$$

From (1),

$$xy = c^2$$

$$\frac{d}{dx}(xy) = \frac{d}{dx}(c^2)$$

$$x \frac{dy}{dx} + y(1) = 0$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

$$\Rightarrow \frac{dy}{dx} = \frac{-y}{x}$$

$$\text{At } P(c, c), \frac{dy}{dx} = \frac{-c}{c} = -1$$

$$y_1 = -1$$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} \left(\frac{dy}{dx} \right) \\ &= \frac{d}{dx} \left(\frac{-y}{x} \right) \\ &= -\left[\frac{x \frac{dy}{dx} - y(1)}{x^2} \right]\end{aligned}$$

$$\therefore \text{At } P(c, c), \frac{d^2y}{dx^2} = -\left[\frac{c(-1) - (c)}{c^2} \right]$$

$$= \frac{2c}{c^2}$$

\therefore

$$y_2 = \frac{2}{c}$$

$$\frac{1+y_1^2}{y_2} = \frac{1+(-1)^2}{\frac{2}{c}} = \frac{2}{\frac{2}{c}} = c$$

Consider y_2

$$\bar{x} = c - (-1)(c) = 2c$$

$$\bar{y} = c + c = 2c$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT 3- DIFFERENTIAL CALCULUS

Circle of Curvature

We know that

$$\rho = \frac{(1+y_1^2)^{3/2}}{y_2}$$

$$\rho = \frac{(1+(-1)^2)^{3/2}}{\frac{2}{c}} = (2)^{3/2} \times \frac{c}{2}$$

∴ at $P(c, c)$,

$$= c\sqrt{2}$$

$$= c\sqrt{2}$$

$$\therefore \rho = c\sqrt{2}$$

∴ (2) becomes

$$(x-2c)^2 + (y-2c)^2 = (c\sqrt{2})^2 .$$