

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

Reduce the quadratic form $\underset{1}{2}x^2 + \underset{2}{2}x^2 + \underset{3}{x^2} + 4x_1x_2 = 0$ to canonical form by orthogonal

reduction .Find rank, index, signature and nature

Step 1:

The matrix form is

$$A = \begin{bmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Step 2:

Characteristic equation ,Eigen values, Eigen vectors

C_1 =Sum of leading diagonal elements

$$=2+2+1=5$$

C_2 = Sum of minors of leading diagonal elements

$$=4$$

$$C_3=|A|$$

$$= \begin{vmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= 0$$

The characteristic equation is

$$\lambda^3 - 5\lambda^2 + 4\lambda = 0$$

The eigen values are 0,1,4

The eigen vectors are $(A - \lambda I)X=0$

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

$$\begin{matrix} 2 & 2 & 0 & 1 & 0 & 0 & x_1 \\ [(2 & 2 & 0) - \lambda(0 & 1 & 0)](x_2) = 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & x_3 \end{matrix}$$

$$\begin{pmatrix} 2-\lambda & 2 & 0 & x_1 \\ 2 & 2-\lambda & 0 & 0 \\ 0 & 0 & 1-\lambda & x_3 \end{pmatrix}(x_2) = 0$$

$$\begin{matrix} 2 & 2 & 0 & x_1 \\ (2 & 2 & 0)(x_2) = 0 \\ 0 & 0 & 1 & x_3 \end{matrix}$$

CASE (i)

When $\lambda = 0$

$$\begin{matrix} 2 & 2 & 0 & x_1 \\ (2 & 2 & 0)(x_2) = 0 \\ 0 & 0 & 1 & x_3 \end{matrix}$$

The cofactor of first row elements are $\begin{pmatrix} 2 & 1 \\ -2 & 0 \end{pmatrix}$ ie (-1)

The Eigen vector when $\lambda = 0$ is $\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$

CASE (ii)

When $\lambda = 1$

$$\begin{matrix} 1 & 2 & 0 & x_1 \\ (2 & 1 & 0)(x_2) = 0 \\ 0 & 0 & 0 & x_3 \end{matrix}$$

The cofactor of third row elements are $\begin{pmatrix} 0 & 0 \\ 0 & -3 \end{pmatrix}$ ie $(0, -1)$

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

The Eigen vector when $\lambda = 1$ is $(\begin{array}{c} 0 \\ 0 \\ -1 \end{array})$

CASE (iii)

When $\lambda = 4$

$$\begin{pmatrix} 1 & 2 & 0 & x_1 \\ 2 & 1 & 0 & x_2 \\ 0 & 0 & 0 & x_3 \end{pmatrix} = 0$$

The cofactor of first row elements are $(6) \text{ ie } (\begin{array}{cc} 6 & 1 \\ 0 & 0 \end{array})$

The Eigen vector when $\lambda = 4$ is $(\begin{array}{c} -2 \\ 1 \\ 1 \end{array})$

STEP 3:

To check pair wise orthogonality

$$X_1^T X_2 = (2 \quad -2 \quad 0) \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} = 0$$

$$X_2^T X_3 = (0 \quad 0 \quad -1) \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = 0$$

$$X_3^T X_1 = (1 \quad 1 \quad 0) \begin{pmatrix} 2 \\ -2 \\ 0 \end{pmatrix} = 0$$

STEP 4:

To find normalized vector

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

Eigen vector	$l(x) = \sqrt{x_1^2 + x_2^2 + x_3^2}$	$x_1/l(x_1)$ Normalized vector $= (x_2/l(x_2))$ $x_3/l(x_3)$
$\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$	$\sqrt{1+1+0} = \sqrt{2}$	$\begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \\ 0 \end{pmatrix}$
$\begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$	$\sqrt{0+0+1} = \sqrt{1}$	$\begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$
$\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$	$\sqrt{1+1+0} = \sqrt{2}$	$\begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{pmatrix}$

STEP 5:

Normalized modal matrix

$$N = \begin{vmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & -1 & 0 \end{vmatrix}$$

$$N^T = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & -1 \\ 1/\sqrt{2} & 1/\sqrt{2} & 1 \end{bmatrix}$$

STEP 6:

$$NN^T = N^TN = I$$

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

$$N^T N = \mathbf{I} \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & -1 \\ h^{1/\sqrt{2}} & h^{1/\sqrt{2}} & h^1 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & -1 & 0 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \mathbf{I}$$

STEP 7:

To find diagonalize matrix

$$N^T A N = D$$

$$N^T A = \mathbf{I} \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 0 & -1 \\ h^{1/\sqrt{2}} & h^{1/\sqrt{2}} & h^1 \end{pmatrix} \begin{pmatrix} 2 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 4/\sqrt{2} & 4/\sqrt{2} & 0 \end{pmatrix}$$

$$N^T A N = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 4/\sqrt{2} & 4/\sqrt{2} & 0 \end{pmatrix} \mathbf{I} \begin{pmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & -1 & 0 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{pmatrix}$$

$$= D$$

UNIT 2 – ORTHOGONAL TRANSFORMATION OF A REAL SYMMETRIC MATRIX

Reduction of quadratic form to canonical form by orthogonal transformation

Step 8:

$$Y^T D Y = 0$$

$$0y_1^2 + y_2^2 + 4y_3^2 = 0$$

The index p=2

Rank r=2

Signature s=2p-r =2

The nature is semi positive