



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

#### JACOBIANS

If  $u = f(x, y)$  &  $v = g(x, y)$  be the two cts. functions of  $x$  &  $y$  then the functional determinant

$$|J| = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} \text{ is called}$$

Jacobians of  $u$  and  $v$  with respect to  $x$  &  $y$ .

Three functions & three variables

$$|J| = \frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix}$$

#### Properties :

1) If  $u, v$  are functions of  $x$  &  $y$  and  $x, y$  are functions of  $r$  &  $s$  then

$$\frac{\partial(u, v)}{\partial(r, s)} = \frac{\partial(u, v)}{\partial(x, y)} \cdot \frac{\partial(x, y)}{\partial(r, s)}$$

2) If  $u$  &  $v$  are functions of  $x$  &  $y$  then

$$\frac{\partial(u, v)}{\partial(x, y)} \cdot \frac{\partial(x, y)}{\partial(u, v)} = 1$$



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

3) If  $u, v, w$  are functionally dependent functions which are depends on  $x, y, z$  then

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = 0$$

① If  $u = \frac{y^2}{x}, v = \frac{x^2}{y}$  Find  $\frac{\partial(u, v)}{\partial(x, y)}$ .

Soln:  $u = \frac{y^2}{x}; v = \frac{x^2}{y}$

$$\frac{\partial u}{\partial x} = -\frac{y^2}{x^2}; \frac{\partial v}{\partial x} = \frac{2x}{y}$$

$$\frac{\partial u}{\partial y} = \frac{2y}{x}; \frac{\partial v}{\partial y} = -\frac{x^2}{y^2}$$

$$|J| = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} = \begin{vmatrix} -y^2/x^2 & 2y/x \\ 2x/y & -x^2/y^2 \end{vmatrix}$$

$$= -\frac{y^2}{x^2} \times -\frac{x^2}{y^2} - \frac{2y}{x} \times \frac{2x}{y}$$

$$= 1 - 4$$

$$= -3$$



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

② If  $u = \frac{yz}{x}$ ,  $v = \frac{zx}{y}$ ,  $w = \frac{xy}{z}$  Find  $\frac{\partial(u,v,w)}{\partial(x,y,z)}$

$$u = \frac{yz}{x} \quad ; \quad v = \frac{zx}{y} \quad ; \quad w = \frac{xy}{z}$$

$$\frac{\partial u}{\partial x} = -\frac{yz}{x^2} \quad ; \quad \frac{\partial v}{\partial x} = \frac{z}{y} \quad ; \quad \frac{\partial w}{\partial x} = \frac{y}{z}$$

$$\frac{\partial u}{\partial y} = \frac{z}{x} \quad ; \quad \frac{\partial v}{\partial y} = -\frac{zx}{y^2} \quad ; \quad \frac{\partial w}{\partial y} = \frac{x}{z}$$

$$\frac{\partial u}{\partial z} = \frac{y}{x} \quad ; \quad \frac{\partial v}{\partial z} = \frac{x}{y} \quad ; \quad \frac{\partial w}{\partial z} = -\frac{xy}{z^2}$$

$$|J| = \frac{\partial(u,v,w)}{\partial(x,y,z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix} = \begin{vmatrix} -\frac{yz}{x^2} & \frac{z}{x} & \frac{y}{z} \\ \frac{z}{y} & -\frac{zx}{y^2} & \frac{x}{z} \\ \frac{y}{z} & \frac{x}{y} & -\frac{xy}{z^2} \end{vmatrix}$$

$$= -\frac{yz}{x^2} \left[ -\frac{zx}{y^2} \times -\frac{xy}{z^2} - \frac{x}{y} \times \frac{x}{z} \right] - \frac{z}{x} \left[ -\frac{xy}{z^2} \times \frac{z}{y} - \frac{x}{y} \times \frac{y}{z} \right] +$$

$$\frac{y}{z} \left[ \frac{z}{y} \times \frac{x}{z} - \frac{y}{z} \times -\frac{zx}{y^2} \right]$$

$$= 4$$



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

③ If  $x = r \cos \theta$ ,  $y = r \sin \theta$  find  $\frac{\partial(x,y)}{\partial(r,\theta)}$

Soln:

$$x = r \cos \theta \quad ; \quad y = r \sin \theta$$

$$\frac{\partial x}{\partial r} = \cos \theta \quad ; \quad \frac{\partial y}{\partial r} = \sin \theta$$

$$\frac{\partial x}{\partial \theta} = -r \sin \theta \quad ; \quad \frac{\partial y}{\partial \theta} = r \cos \theta$$

$$|J| = \frac{\partial(x,y)}{\partial(r,\theta)} = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix}$$

$$= r \cos^2 \theta + r \sin^2 \theta = r$$

⑦ If  $x = uv$ ,  $y = \frac{u+v}{u-v}$  find  $\frac{\partial(u,v)}{\partial(x,y)}$

Soln:

Wkt  $\frac{\partial(x,y)}{\partial(u,v)} \times \frac{\partial(u,v)}{\partial(x,y)} = 1$

$$\frac{\partial(x,y)}{\partial(u,v)} + \frac{\partial(u,v)}{\partial(x,y)} = 1$$

$$J \cdot J' = 1$$

$$|J| = \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

$$\begin{aligned}x &= uv & ; & \quad y = \frac{u+v}{u-v} \\ \frac{\partial x}{\partial u} &= v & \quad \frac{\partial y}{\partial u} &= \frac{(u-v) - (u+v)}{(u-v)^2} = \frac{-2v}{(u-v)^2} \\ \frac{\partial x}{\partial v} &= u & \quad \frac{\partial y}{\partial v} &= \frac{(u-v) - (u+v)(-1)}{(u-v)^2} = \frac{2u}{(u-v)^2}\end{aligned}$$

$$\therefore |J| = \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} v & u \\ \frac{-2v}{(u-v)^2} & \frac{2u}{(u-v)^2} \end{vmatrix}$$

$$= \frac{2uv}{(u-v)^2} + \frac{2uv}{(u-v)^2} = \frac{4uv}{(u-v)^2}$$

$$\frac{\partial(x,y)}{\partial(u,v)} \times \frac{\partial(u,v)}{\partial(x,y)} = 1$$

$$\begin{aligned}\frac{\partial(u,v)}{\partial(x,y)} &= \frac{1}{\frac{\partial(x,y)}{\partial(u,v)}} = \frac{1}{\frac{4uv}{(u-v)^2}} \\ &= \frac{(u-v)^2}{4uv}\end{aligned}$$

Q) If  $x = r \cos \alpha$ ,  $y = r \sin \alpha$  find  $\frac{\partial(x,y)}{\partial(r,\alpha)}$





## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

If  $u = 2xy$ ,  $v = x^2 - y^2$  and  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

find  $\frac{\partial(u,v)}{\partial(r,\theta)}$

Soln:  $\frac{\partial(u,v)}{\partial(r,\theta)} = \frac{\partial(u,v)}{\partial(x,y)} \cdot \frac{\partial(x,y)}{\partial(r,\theta)}$

Eqn:  $u = 2xy$        $v = x^2 - y^2$

$$\frac{\partial u}{\partial x} = 2y$$

$$\frac{\partial v}{\partial x} = 2x$$

$$\frac{\partial u}{\partial y} = 2x$$

$$\frac{\partial v}{\partial y} = -2y$$

$$\frac{\partial(u,v)}{\partial(x,y)} = \begin{vmatrix} 2y & 2x \\ 2x & -2y \end{vmatrix} = -4y^2 - 4x^2$$

Eqn:  $x = r \cos \theta$        $y = r \sin \theta$

$$\frac{\partial x}{\partial r} = \cos \theta$$

$$\frac{\partial y}{\partial r} = \sin \theta$$

$$\frac{\partial x}{\partial \theta} = -r \sin \theta$$

$$\frac{\partial y}{\partial \theta} = r \cos \theta$$

$$\frac{\partial(x,y)}{\partial(r,\theta)} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r \cos^2 \theta + r \sin^2 \theta = r$$



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

$$\begin{aligned} \frac{\partial(u,v)}{\partial(x,y)} &= (-4y^2 - 4x^2) \times x \\ &= -4(x^2 + y^2) \times x \\ &= -4x^2 \times x \\ &= -4x^3 \end{aligned}$$

$$x = r \cos \theta, y = r \sin \theta$$

$$x^2 + y^2 = r^2 \cos^2 \theta + r^2 \sin^2 \theta$$

$$= r^2$$

10) S.T. the functions  $u = \frac{x}{y}$  &  $v = \frac{x+y}{x-y}$  are functionally dependent and find the relationship btwn. them.

Soln:  $u = \frac{x}{y}$

$v = \frac{x+y}{x-y}$

$$\frac{\partial u}{\partial x} = \frac{1}{y}$$

$$\frac{\partial v}{\partial x} = \frac{x-y - (x+y)(-1)}{(x-y)^2} = \frac{-2xy}{(x-y)^2}$$

$$\frac{\partial u}{\partial y} = -\frac{x}{y^2}$$

$$\frac{\partial v}{\partial y} = \frac{x-y - (x+y)(-1)}{(x-y)^2} = \frac{2x}{(x-y)^2}$$

$$\frac{\partial(u,v)}{\partial(x,y)} = \begin{vmatrix} \frac{1}{y} & -\frac{x}{y^2} \\ -\frac{2xy}{(x-y)^2} & \frac{2x}{(x-y)^2} \end{vmatrix}$$

$$= 0$$

$\therefore$  the .eqn. functions are functionally dependent



## DEPARTMENT OF MATHEMATICS

### UNIT - IV FUNCTIONS OF SEVERAL VARIABLES

Ex:  $u = \frac{x}{y}, v = \frac{x+y}{x-y}$

$$v = \frac{\left[\frac{x}{y} + 1\right]y}{\left[\frac{x}{y} - 1\right]y} = \frac{u+1}{u-1}$$

$$\Rightarrow v = \frac{u+1}{u-1}$$

11) Ex: The functions  $u = 2x - y + 3z, v = 2x - y - z, w = 2x - y + z$  are functionally dependent. Find relationship between them.

Ans: functionally dependent.

Relationship:  $u + v = 2w$ .

12) Ex: If  $u = xy + yz + zx, v = x^2 + y^2 + z^2$  &  $w = x + y + z$  determine whether a functional relation between  $x, y, z$  are dependent & find relationship between them.

Ans:  $2u + v = w^2$  (relationship)