



DEPARTMENT OF MATHEMATICS

Moment Generating function :

The moment generating function (m.g.f) of a random variable 'x' (about origin) whose probability function is given by,

$$M_x(t) = E[e^{tx}]$$

For a discrete random variable, m.g.f is given by

$$M_x(t) = \sum e^{tx} P(x)$$

For a continuous random variable, m.g.f is given by

$$M_x(t) = \int_{-\infty}^{\infty} e^{tx} f(x) dx$$

Problems :

- ① Prove that the r^{th} moment of the r.v 'x' about origin is $M_x(t) = \sum_{r=0}^{\infty} \frac{t^r}{r!} \mu_r'$

Solution :

$$M_x(t) = E(e^{tx})$$

$$= E \left[1 + \frac{tx}{1!} + \frac{t^2 x^2}{2!} + \dots + \frac{t^r x^r}{r!} + \dots \right]$$

$$= E(1) + E \left[\frac{tx}{1!} \right] + E \left[\frac{t^2 x^2}{2!} \right] + \dots + E \left[\frac{t^r x^r}{r!} \right] + \dots$$

$$= 1 + t E[X] + \frac{t^2}{2!} E(X^2) + \dots +$$

$$= 1 + t \mu_1' + \frac{t^2}{2!} \mu_2' + \dots + \frac{t^r}{r!} \mu_r' + \dots$$

$$M_x(t) = \sum_{r=0}^{\infty} \frac{t^r}{r!} \mu_r'$$



DEPARTMENT OF MATHEMATICS

② A random variable X has the probability function...

$$f(x) = \frac{1}{2^x}, \quad x = 1, 2, 3, \dots, \infty$$

Find its (i) M.G.F (ii) Mean & Variance (iii) $P(X \text{ is even})$
Solution: (iv) $P(X \geq 5)$ and $\{P(X \text{ is divisible by } 3)\}$

(i) M.G.F = $M_X(t) = E(e^{tx})$

$$= \sum_{x=1}^{\infty} e^{tx} f(x)$$

$$= \sum_{x=1}^{\infty} e^{tx} \cdot \frac{1}{2^x}$$

$$= \sum_{x=1}^{\infty} \left(\frac{e^t}{2}\right)^x$$

$$= \frac{e^t}{2} + \left(\frac{e^t}{2}\right)^2 + \left(\frac{e^t}{2}\right)^3 + \dots$$

$$= \frac{e^t}{2} \left[1 + \frac{e^t}{2} + \left(\frac{e^t}{2}\right)^2 + \dots\right]$$

$$= \frac{e^t}{2} \left(1 - \frac{e^t}{2}\right)^{-1}$$

$$= \frac{e^t}{2} \cdot \frac{2}{2 - e^t}$$

$$M_X(t) = \frac{e^t}{2 - e^t}$$

(ii) Mean = $E(X) = M_X'(0)$

$$M_X'(t) = \frac{(2 - e^t)e^t - e^t(-e^t)}{(2 - e^t)^2}$$

$$= \frac{2e^t - e^{2t} + e^{2t}}{(2 - e^t)^2}$$

$$M_X'(t) = \frac{2e^t}{(2 - e^t)^2}$$



DEPARTMENT OF MATHEMATICS

$$E(X) = M_x'(0) = \frac{2}{2^2 - 1} = \frac{1}{2} \cdot 2$$

$$\boxed{E(X) = \frac{1}{2}}$$

$$M_x''(t) = \frac{(2 - e^t)^2 (2e^t) - 2e^t \cdot 2(2 - e^t)(-e^t)}{(2 - e^t)^4}$$

$$= \frac{2e^t(2 - e^t) [2 - e^t + 2e^t]}{(2 - e^t)^4}$$

$$= \frac{2e^t(2 + e^t)}{(2 - e^t)^3}$$

$$M_x''(0) = \frac{2(2+1)}{(2-1)^3} = 2(3) = 6$$

$$\therefore \underline{E(X^2) = 6}$$

$$\begin{aligned} \therefore \text{Variance} &= E(X^2) - [E(X)]^2 \\ &= 6 - 2^2 \\ &= 6 - 4 \end{aligned}$$

$$\boxed{\text{Variance} = 2}$$

$$(iii) P(X \text{ is even}) = P(X=2) + P(X=4) + \dots$$

$$= \frac{1}{2^2} + \frac{1}{2^4} + \dots$$

$$= \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^4 + \dots$$

$$= \frac{\left(\frac{1}{2}\right)^2}{1 - \left(\frac{1}{2}\right)^2}$$

$$= \frac{1/4}{1 - 1/4}$$

$$= \frac{1}{2}$$

[\because It is a geometric series = $\frac{a}{1-r}$]

$a \rightarrow$ first term

$r \rightarrow$ Common ratio



DEPARTMENT OF MATHEMATICS

$$\begin{aligned} \text{(iv) } P(X \geq 5) &= P(X=5) + P(X=6) + P(X=7) + \dots \infty \\ &= \frac{1}{2^5} + \frac{1}{2^6} + \frac{1}{2^7} + \dots \infty \\ &= \left(\frac{1}{2}\right)^5 + \left(\frac{1}{2}\right)^6 + \left(\frac{1}{2}\right)^7 + \dots \infty \\ &= \frac{\left(\frac{1}{2}\right)^5}{1 - \frac{1}{2}} = \frac{\frac{1}{32}}{\frac{1}{2}} = \frac{1}{16} \end{aligned}$$

$$P(X \geq 5) = \frac{1}{16}$$

$$\begin{aligned} P(X \text{ is divisible by } 3) &= P(X=3) + P(X=6) + \dots \infty \\ &= \left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^6 + \dots \infty \\ &= \frac{\left(\frac{1}{2}\right)^3}{1 - \left(\frac{1}{2}\right)^3} = \frac{\frac{1}{8}}{\frac{7}{8}} = \frac{1}{7} \end{aligned}$$

$$P(X \text{ is divisible by } 3) = \frac{1}{7}$$

③ Find the MGF for the distribution where

$$f(x) = \begin{cases} \frac{2}{3} & \text{at } x=1 \\ \frac{1}{3} & \text{at } x=2 \\ 0 & \text{otherwise} \end{cases}$$

Solution :

$$\begin{aligned} M_X(t) &= E(e^{tx}) \\ &= \sum_{x=0}^{\infty} e^{tx} f(x) \\ &= e^0 f(0) + e^t f(1) + e^{2t} f(2) + \dots \infty \\ &= 0 + e^t \cdot \frac{2}{3} + e^{2t} \left(\frac{1}{3}\right) + 0 : \end{aligned}$$

$$M_X(t) = \frac{2e^t}{3} + \frac{e^{2t}}{3}$$



DEPARTMENT OF MATHEMATICS

④ Find the MGF of a random variable 'x' having the density function

$$f(x) = \begin{cases} \frac{x}{2}, & 0 \leq x \leq 2 \\ 0, & \text{otherwise} \end{cases}$$

Using the generating fn find the first four moments about the origin.

Solution:

(i) $M_x(t) = \int_{-\infty}^{\infty} e^{tx} f(x) dx$

$$= \int_0^2 e^{tx} \cdot \frac{x}{2} dx$$

$$= \frac{1}{2} \int_0^2 x e^{tx} dx$$

$$= \frac{1}{2} \left[\frac{x e^{tx}}{t} - \frac{e^{tx}}{t^2} \right]_0^2$$

$$= \frac{1}{2} \left[\frac{2e^{2t}}{t} - \frac{e^{2t}}{t^2} + \frac{1}{t^2} \right]$$

$$M_x(t) = \frac{1}{2t^2} [2te^{2t} - e^{2t} + 1]$$

$$u = x, v = e^t$$

$$u' = 1, v' = \frac{e^t}{t}$$

$$v_2 = \frac{e^t}{t^2}$$

$$u v_1 - u' v_2$$

(ii) $M_x^2(t) = \frac{2te^{2t}}{2t^2} - \frac{e^{2t}}{2t^2} + \frac{1}{2t^2}$

$$= t^{-1} e^{2t} - \frac{1}{2} t^{-2} e^{2t} + \frac{1}{2} t^{-2}$$

$$M_x'(t) = e^{2t} \left(\frac{-1}{t^2} \right) + t^{-1} \cdot \frac{e^{2t}}{2} - \frac{1}{2} \left[\frac{e^{2t}}{2} t^{-2} + \frac{e^{2t}}{t^3} \right]$$

$$+ \frac{1}{2} (-2) \frac{1}{t^3}$$

$$= -\frac{e^{2t}}{t^2} + \frac{e^{2t}}{2t} - \frac{e^{2t}}{4t^2} + \frac{e^{2t}}{t^3} - \frac{1}{t^3}$$



DEPARTMENT OF MATHEMATICS

⑤ Let 'x' be a random variable with p.d.f

$$f(x) = \begin{cases} \frac{1}{3} e^{-x/3}, & x > 0 \\ 0, & \text{otherwise} \end{cases}$$

Find (i) $P(x > 3)$ (ii) MGF of x (iii) $E(x)$ and $\text{Var}(x)$.

Solution:
Given: $f(x) = \begin{cases} \frac{1}{3} e^{-x/3}, & x > 0 \\ 0, & \text{otherwise.} \end{cases}$

$$\begin{aligned} \text{(i) } P(x > 3) &= \int_3^{\infty} f(x) dx \\ &= \int_3^{\infty} \frac{1}{3} e^{-x/3} dx \\ &= \frac{1}{3} \left[\frac{e^{-x/3}}{-1/3} \right]_3^{\infty} \\ &= e^{-1} \end{aligned}$$

$$P(x > 3) = 0.3679$$

(ii) MGF of x:

$$\begin{aligned} M_x(t) &= E(e^{tx}) \\ &= \int_{-\infty}^{\infty} e^{tx} f(x) dx \\ &= \int_0^{\infty} e^{tx} \frac{1}{3} e^{-x/3} dx \\ &= \frac{1}{3} \int_0^{\infty} e^{(t - \frac{1}{3})x} dx = \frac{1}{3} \int_0^{\infty} e^{-\left(\frac{1}{3} - t\right)x} dx \\ &= \frac{1}{3} \left[\frac{e^{-\left(\frac{1}{3} - t\right)x}}{-\left(\frac{1}{3} - t\right)} \right]_0^{\infty} \end{aligned}$$



DEPARTMENT OF MATHEMATICS

$$M_x(t) = -\frac{1}{3} \left[0 - \frac{1}{\frac{1}{3} - t} \right] = +\frac{1}{3} \left(\frac{1}{\frac{1}{3} - t} \right)$$

$$\boxed{M_x(t) = \frac{1}{1-3t}}$$

$$(iii) \quad M_x(t) = \frac{1}{1-3t} = (1-3t)^{-1}$$

$$M_x'(t) = -\frac{1}{(1-3t)^2} (-3) = \frac{3}{(1-3t)^2}$$

$$M_x'(0) = \frac{3}{1-0} = 3$$

$$\boxed{E(x) = \text{Mean} = M_x'(0) = 3}$$

$$M_x''(t) = -6(1-3t)^{-3} (-3) \\ = 18(1-3t)^{-3}$$

$$M_x''(0) = 18$$

$$E(x^2) = M_x''(0) = 18$$

$$\text{Var}(x) = E(x^2) - [E(x)]^2 \\ = 18 - (3)^2 \\ = 18 - 9$$

$$\boxed{\text{Var}(x) = 9}$$