



DEPARTMENT OF MATHEMATICS

DISCRETE DISTRIBUTIONS :

The important discrete distributions of a random variable 'x' are,

1. Binomial distribution
2. Poisson distribution
3. Geometric distribution
4. Negative Binomial distribution.

BINOMIAL DISTRIBUTION :

The probability mass function of a random variable 'x' which follows the binomial distribution is,

$$P(X=x) = {}^n C_x p^x q^{n-x}, \quad x = 0, 1, 2, \dots, n \quad \&$$

$$(q+p)^n = q^n + {}^n C_1 q^{n-1} p^1 + {}^n C_2 q^{n-2} p^2 + \dots + {}^n C_x p^x q^{n-x}$$

$p+q=1$

which is a binomial series and hence the distribution is called a Binomial distribution.

NOTE :

$$P(0 \text{ Success}) = {}^n C_0 p^0 q^{n-0} = q^n$$

$$P(1 \text{ Success}) = {}^n C_1 p^1 q^{n-1}$$

$$P(2 \text{ Success}) = {}^n C_2 p^2 q^{n-2} \quad \text{and so on.}$$

ASSUMPTIONS :

- (i) There are only two possible outcomes for each trial (Success or failure)
- (ii) The probability of a success is the same for each trial
- (iii) There are 'n' trials, where 'n' is a constant.
- (iv) The 'n' trials are independent.



DEPARTMENT OF MATHEMATICS

Moment Generating function (M.G.F) :

$$\begin{aligned}
 M_x(t) &= E(e^{tx}) \\
 &= \sum_{x=0}^n e^{tx} p(x) \\
 &= \sum_{x=0}^n e^{tx} nC_x p^x q^{n-x} \\
 &= \sum_{x=0}^n (pe^t)^x nC_x q^{n-x} \\
 &= q^n + nC_1 (pe^t) q^{n-1} + nC_2 (pe^t)^2 q^{n-2} + \dots
 \end{aligned}$$

$$M_x(t) = (q + pe^t)^n$$

$$\begin{aligned}
 (x+a)^n &= a^n + nC_1 x a^{n-1} \\
 &+ nC_2 x^2 a^{n-2} + \dots \\
 &+ nC_n x^n a
 \end{aligned}$$

Mean and Variance :

$$M_x(t) = (q + pe^t)^n$$

$$M_x'(t) = n(q + pe^t)^{n-1} pe^t$$

$$M_x'(0) = n(q + p)^{n-1} p$$

$$M_x'(0) = np$$

$$(\because p + q = 1)$$

$$\therefore \text{Mean} = E(x) = np$$



DEPARTMENT OF MATHEMATICS

$$M_x''(t) = np \left[(\alpha + pe^t)^{n-1} \cdot e^t + e^t (n-1)(\alpha + pe^t)^{n-2} pe^t \right]$$

Putting $t=0$,

$$M_x''(0) = np \left[(\alpha + p)^{n-1} + (n-1)(\alpha + p)^{n-2} p \right]$$

$$= np \left[1 + (n-1)p \right]$$

$$= np + n^2 p^2 - np^2$$

$$M_x''(0) = n^2 p^2 + np(1-p)$$

$$\boxed{M_x''(0) = E(x^2) = n^2 p^2 + np\alpha}$$

$$\therefore \text{Variance} = E(x^2) - [E(x)]^2$$

$$= n^2 p^2 + np\alpha - (np)^2$$

$$= n^2 p^2 + np\alpha - n^2 p^2$$

$$\boxed{\text{Variance} = np\alpha}$$

$$\text{Standard deviation} = \sqrt{\text{Variance}}$$

$$\boxed{SD = \sqrt{np\alpha}}$$



DEPARTMENT OF MATHEMATICS

PROBLEMS :

- ① The mean and SD of a binomial distribution are 20 and 4. Determine the distribution.

Solution :

$$\text{Mean} = np = 20 \rightarrow \textcircled{1}$$

$$\text{SD} = \sqrt{npq} = 4$$

$$npq = 4^2 = 16 \rightarrow \textcircled{2}$$

$$\frac{\textcircled{2}}{\textcircled{1}} = \frac{npq}{np} = \frac{16}{20} \Rightarrow \boxed{q = \frac{4}{5}}$$

$$p = 1 - q = 1 - \frac{4}{5} = \frac{1}{5}$$

$$\boxed{p = \frac{1}{5}}$$

subs p in $\textcircled{1}$,

$$n \cdot \frac{1}{5} = 20$$

$$\boxed{n = 100}$$

\therefore The binomial distribution is,

$$P(X = x) = P(x) = {}^n C_x p^x q^{n-x} \\ = {}^{100} C_x \left(\frac{1}{5}\right)^x \left(\frac{4}{5}\right)^{100-x}, x = 0, 1, \dots, 100$$

- ② If X is a binomial random variable with expected value 2 and variance $4/3$, find $P(X=5)$.

Solution :

$$E(X) = 2$$

$$np = 2 \rightarrow \textcircled{1}$$

$$\text{variance} = 4/3$$

$$npq = 4/3 \rightarrow \textcircled{2}$$



DEPARTMENT OF MATHEMATICS

$$\frac{(2)}{(1)} = \frac{npq}{np} = \frac{4/3}{2} = \frac{4}{3} \times \frac{1}{2} = \frac{2}{3}$$

$$q = \frac{2}{3}$$

$$p = 1 - q = 1 - \frac{2}{3} = \frac{1}{3} \Rightarrow p = \frac{1}{3}$$

subs p in (1),

$$n \cdot \frac{1}{3} = 2 \Rightarrow n = 6$$

$$P(X=x) = {}^n C_x p^x q^{n-x}$$

$$P(X=5) = {}^6 C_5 \left(\frac{1}{3}\right)^5 \left(\frac{2}{3}\right)^{6-5}$$

$$= {}^6 C_1 \frac{1}{3^5} \cdot \frac{2}{3} = \frac{6 \times 2}{3 \times 3 \times 3 \times 3 \times 3}$$

$$P(X=5) = 0.0165$$

- (3) If the MGF of a r.v x is of the form $(0.4e^t + 0.6)^8$.
 What is the MGF of $3x+2$. Find $E(x)$, $\text{Var}(x)$, $P(X=2)$.

Solution:

$$M_x(t) = (0.4e^t + 0.6)^8 = E(e^{tx})$$

$$\text{M.G.F of } 3x+2 = E(e^{t(3x+2)})$$

$$= E(e^{2t} \cdot e^{3tx})$$

$$= e^{2t} E(e^{3tx})$$

$$= e^{2t} E(e^{x(3t)})$$

$$M_{3x+2}(t) = e^{2t} (0.4e^{3t} + 0.6)^8$$

$$M_x(t) = (0.4e^t + 0.6)^8$$

$$\text{But } M_x(t) = (q + pe^t)^n$$

$$\therefore q = 0.6, p = 0.4 \text{ \& } n = 8$$



DEPARTMENT OF MATHEMATICS

$$E(X) = np = 8 \times 0.4 = 3.2$$

$$E(X) = 3.2$$

$$\text{Variance} = E(X^2) - n p q = 8 \times 0.4 \times 0.6$$

$$\text{Variance} = 1.92$$

$$P(X=x) = {}^n C_x p^x q^{n-x}$$

$$P(X=2) = {}^8 C_2 (0.4)^2 (0.6)^{8-2}$$
$$= \frac{8 \times 7}{2} \times 0.16 \times 0.0467$$

$$P(X=2) = 0.2092$$

- 4) If 10% of the screws produced by an automatic machine are defective, find the probability that out of 20 screws selected at random, there are (i) exactly 2 defective (ii) at most 3 defective (iii) at least 2 defectives and (iv) between 1 and 3 defectives (inclusive).

Solution:

Let x be the R.V denoting the number of defective screws.

$$p = 10\% = \frac{10}{100} = \frac{1}{10}$$

$$q = 1 - p = 1 - \frac{1}{10} = \frac{9}{10}$$

$$n = 20$$

$$P(X=x) = {}^n C_x p^x q^{n-x}$$

(i) $P(\text{getting exactly 2 defectives})$

$$= P(X=2)$$

$$= {}^{20} C_2 \left(\frac{1}{10}\right)^2 \left(\frac{9}{10}\right)^{20-2}$$

$$= \frac{20 \times 19}{2} \times \frac{1}{100} \times \left(\frac{9}{10}\right)^{18}$$



DEPARTMENT OF MATHEMATICS

$$P(\text{getting exactly 2 defectives}) = \underline{0.2852}$$

$$(ii) P(\text{getting atmost 3 defective}) = P(X \leq 3)$$

$$= P(X=0) + P(X=1) + P(X=2) + P(X=3)$$

$$= {}^{20}C_0 \left(\frac{1}{10}\right)^0 \left(\frac{9}{10}\right)^{20-0} + {}^{20}C_1 \left(\frac{1}{10}\right)^1 \left(\frac{9}{10}\right)^{20-1} +$$
$${}^{20}C_2 \left(\frac{1}{10}\right)^2 \left(\frac{9}{10}\right)^{20-2} + {}^{20}C_3 \left(\frac{1}{10}\right)^3 \left(\frac{9}{10}\right)^{20-3}$$

$$= \left(\frac{9}{10}\right)^{20} + \frac{20}{10} \left(\frac{9}{10}\right)^{19} + \frac{20 \times 19}{2} \times \frac{1}{100} \left(\frac{9}{10}\right)^{18} +$$

$$\frac{20 \times 19 \times 18}{2 \times 3} \times \frac{1}{1000} \left(\frac{9}{10}\right)$$

$$= 0.1216 + 0.2702 + 0.2852 + 0.1901$$

$$= \underline{0.8671}$$

$$(iii) P(\text{getting atleast 2 defective}) = P(X \geq 2)$$

$$= 1 - P(X < 2)$$

$$= 1 - [P(X=0) + P(X=1) + P(X=2)]$$

$$= 1 - [0.1216 + 0.2702 + 0.2852]$$

$$= \underline{0.323}$$

$$(iv) P(\text{getting number of defectives between 1 and 3})$$

$$= P(1 \leq X \leq 3) \quad (\text{including})$$

$$= P(X=1) + P(X=2) + P(X=3)$$

$$= 0.2702 + 0.2852 + 0.1901$$

$$= \underline{0.7455}$$