



**DEPARTMENT OF MATHEMATICS**

INVERSE OF A MATRIX - GAUSS JORDAN METHOD

36

Let us find the inverse of a non-singular square matrix  $A$  of order three. If  $X$  is the inverse of  $A$ , then  $AX = I$  where  $I$  is the unit matrix of order 3. Now, we have to find the elements of  $X$ .

$$\text{Let } A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \text{ and } X = \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix}$$

Therefore  $AX = I$  reduces to,

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \rightarrow \textcircled{1}$$

This equation is equivalent to the three equations given below:

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} x_{11} \\ x_{21} \\ x_{31} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad \rightarrow \textcircled{2}$$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} x_{12} \\ x_{22} \\ x_{32} \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \rightarrow \textcircled{3}$$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} x_{13} \\ x_{23} \\ x_{33} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad \rightarrow \textcircled{4}$$



**DEPARTMENT OF MATHEMATICS**

From equations (2), (3), (4), we can solve for the vectors  $\begin{pmatrix} x_{11} \\ x_{21} \\ x_{31} \end{pmatrix}$ ,  $\begin{pmatrix} x_{12} \\ x_{22} \\ x_{32} \end{pmatrix}$  and  $\begin{pmatrix} x_{13} \\ x_{23} \\ x_{33} \end{pmatrix}$  by Gauss-Jordan method. The solution set of each system (2), (3), (4) will be the corresponding column of the inverse matrix  $X$ .

Problems:

- (1) Find the inverse of the matrix  $\begin{pmatrix} 5 & -2 \\ 3 & 4 \end{pmatrix}$  by Gauss-Jordan method.

Solution:

$$\text{Let } A = \begin{pmatrix} 5 & -2 \\ 3 & 4 \end{pmatrix} \text{ and } X = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} \text{ be}$$

the inverse of  $A$ , so that  $AX = I$ .

Step 1: The augmented matrix is,

$$\left[ \begin{array}{cc|cc} 5 & -2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right]$$

Step 2:  $R_1 \rightarrow R_1 / 5$

$$\left[ \begin{array}{cc|cc} 1 & -2/5 & 1/5 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right]$$

Step 3:  $R_2 \rightarrow R_2 - 3R_1$

$$\left[ \begin{array}{cc|cc} 1 & -2/5 & 1/5 & 0 \\ 0 & 26/5 & -3/5 & 1 \end{array} \right]$$



**DEPARTMENT OF MATHEMATICS**

Step 4:  $R_2 \rightarrow R_2 / 26/5$

$$\left[ \begin{array}{cc|cc} 1 & -2/5 & 1/5 & 0 \\ 0 & 1 & -3/26 & 5/26 \end{array} \right]$$

Step 5:  $R_1 \rightarrow R_1 + (2/5) R_2$

$$\left[ \begin{array}{cc|cc} 1 & 0 & 2/13 & 1/13 \\ 0 & 1 & -3/26 & 5/26 \end{array} \right]$$

Hence the inverse of the given matrix is,

$$\begin{bmatrix} 2/13 & 1/13 \\ -3/26 & 5/26 \end{bmatrix} = \frac{1}{26} \begin{bmatrix} 4 & 2 \\ -3 & 5 \end{bmatrix}$$

② Find the inverse of  $\begin{pmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{pmatrix}$  using Gauss-Jordan method.

Solution:

Let  $A = \begin{pmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{pmatrix}$  and  $X = \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix}$

be the inverse of  $A$ , so that  $AX = I$ .

Step 1: The augmented matrix is,

$$\left( \begin{array}{ccc|ccc} 1 & 1 & 3 & 1 & 0 & 0 \\ 1 & 3 & -3 & 0 & 1 & 0 \\ -2 & -4 & -4 & 0 & 0 & 1 \end{array} \right)$$

Step 2:  $R_2 \rightarrow R_2 - R_1$ ,  $R_3 \rightarrow R_3 + 2R_1$

$$\left( \begin{array}{ccc|ccc} 1 & 1 & 3 & 1 & 0 & 0 \\ 0 & 2 & -6 & -1 & 1 & 0 \\ 0 & -2 & 2 & 2 & 0 & 1 \end{array} \right)$$



**DEPARTMENT OF MATHEMATICS**

Step 3:  $R_3 \rightarrow R_3 + R_2$

$$\left( \begin{array}{ccc|ccc} 1 & 1 & 3 & 1 & 0 & 0 \\ 0 & 2 & -6 & -1 & 1 & 0 \\ 0 & 0 & -4 & 1 & 1 & 1 \end{array} \right)$$

Step 4:  $R_1 \rightarrow 2R_1 + R_2$

$$\left( \begin{array}{ccc|ccc} 2 & 4 & 0 & 1 & 1 & 0 \\ 0 & 2 & -6 & -1 & 1 & 0 \\ 0 & 0 & -4 & 1 & 1 & 1 \end{array} \right)$$

Step 5:  $R_2 \rightarrow -4R_2 + 6R_3$

$$\left( \begin{array}{ccc|ccc} 2 & 4 & 0 & 1 & 1 & 0 \\ 0 & -8 & 0 & 10 & 2 & 6 \\ 0 & 0 & -4 & 1 & 1 & 1 \end{array} \right)$$

Step 6:  $R_1 \rightarrow 2R_1 + R_2$

$$\left( \begin{array}{ccc|ccc} 4 & 0 & 0 & 12 & 4 & 6 \\ 0 & -8 & 0 & 10 & 2 & 6 \\ 0 & 0 & -4 & 1 & 1 & 1 \end{array} \right)$$

Step 7:  $R_1 / 4, R_2 / (-8), R_3 / -4$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & 1 & 3/2 \\ 0 & 1 & 0 & -5/4 & -1/4 & -3/4 \\ 0 & 0 & 1 & -1/4 & -1/4 & -1/4 \end{array} \right)$$

Hence the inverse of the given matrix is

$$\begin{pmatrix} 3 & 1 & 3/2 \\ -5/4 & -1/4 & -3/4 \\ -1/4 & -1/4 & -1/4 \end{pmatrix}$$