



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

Unit-II

vector calculus

Gradient:

Let  $\phi(x, y, z)$  be a scalar point function and is continuously differentiable. Then the vector

$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$  is called the gradient of the scalar fn.  $\phi$ .

$$\text{ie., } \text{grad } \phi = \nabla \phi$$

Problems

Q1 Find  $\nabla \phi$  where  $\phi = x^2 + y^2 + z^2$

Soln.

$$\text{Grad. } \phi \text{ (or) } \nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$= \vec{i} \frac{\partial (x^2 + y^2 + z^2)}{\partial x} + \vec{j} \frac{\partial (x^2 + y^2 + z^2)}{\partial y} + \vec{k} \frac{\partial (x^2 + y^2 + z^2)}{\partial z}$$

$$= \vec{i}(2x) + \vec{j}(2y) + \vec{k}(2z)$$

$$\nabla \phi = 2x\vec{i} + 2y\vec{j} + 2z\vec{k}$$

Q2. Find  $\nabla \phi$  where  $\phi = 3x^2y - y^3z^2$  at  $(1, 1, 1)$

Soln.

$$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$= \vec{i} \frac{\partial (3x^2y - y^3z^2)}{\partial x} + \vec{j} \frac{\partial (3x^2y - y^3z^2)}{\partial y} + \vec{k} \frac{\partial (3x^2y - y^3z^2)}{\partial z}$$

$$= \vec{i}[6xy] + \vec{j}[3x^2 - 3y^2z^2] + \vec{k}[0 - 2y^3z]$$

$$\nabla \phi = 6xy\vec{i} + (3x^2 - 3y^2z^2)\vec{j} - 2y^3z\vec{k}$$

$$\nabla \phi_{(1, 1, 1)} = 6(1)(1)\vec{i} + (3 - 3)\vec{j} - 2(1)(1)\vec{k}$$

$$= 6\vec{i} + 0\vec{j} - 2\vec{k}$$

$$= 6\vec{i} - 2\vec{k}$$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

- 3]. Find the maximum directional derivative  
of  $\phi = xyz^2$  at  $(1, 0, 3)$

Soln.

$$\begin{aligned}\nabla\phi &= \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z} \\ &= \vec{i} \frac{\partial}{\partial x} (xyz^2) + \vec{j} \frac{\partial}{\partial y} (xyz^2) + \vec{k} \frac{\partial}{\partial z} (xyz^2) \\ \nabla\phi &= \vec{i}(yz^2) + \vec{j}(xz^2) + \vec{k}(xy^2) \\ \nabla\phi_{(1, 0, 3)} &= \vec{i}(0) + \vec{j}(1)(9) + \vec{k}(0) \\ &= 9\vec{j} \quad \text{maximum DD} = \sqrt{81} = 9\end{aligned}$$

- 4]. Find  $\nabla\phi$  where  $\phi = xyz$  at  $(1, 2, 3)$

Soln.

$$\begin{aligned}\nabla\phi &= \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z} \\ &= \vec{i} \frac{\partial(xyz)}{\partial x} + \vec{j} \frac{\partial(xyz)}{\partial y} + \vec{k} \frac{\partial(xyz)}{\partial z} \\ \nabla\phi &= \vec{i}(yz) + \vec{j}(xz) + \vec{k}(xy) \\ \nabla\phi_{(1, 2, 3)} &= \vec{i}(2)(3) + \vec{j}(1)(3) + \vec{k}(1)(2) \\ &= 6\vec{i} + 3\vec{j} + 2\vec{k}\end{aligned}$$

- 5]. If  $\nabla\phi = yz\vec{i} + zx\vec{j} + xy\vec{k}$ , find  $\phi$ .

Soln.

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$\nabla\phi = \vec{i}(yz) + \vec{j}(zx) + \vec{k}(xy)$$

Equating w.r.t to  $\vec{i}, \vec{j}, \vec{k}$

$$\begin{array}{l|l|l} \frac{\partial\phi}{\partial x} = yz & \frac{\partial\phi}{\partial y} = zx & \frac{\partial\phi}{\partial z} = xy \\ \text{Integrate w.r.t to } x & \text{w.r.t to } y & \text{w.r.t to } z \\ \phi = xyz + f(y, z) & \phi = xyz + g(x, z) & \phi = xyz + h(x, y) \end{array}$$

In general,

$$\phi = xyz + C$$



Scanned with CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

Q. If  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$ , such that  $|\vec{r}| = r$ ,

Prove that

$$\text{i)} \quad \nabla r = \frac{\vec{r}}{r} = \hat{r}$$

$$\text{iii)} \quad \nabla r^n = n r^{n-2} \vec{r}$$

$$\text{ii)} \quad \nabla \left( \frac{1}{r} \right) = - \frac{\vec{r}}{r^3} = - \frac{\hat{r}}{r^2}$$

$$\text{iv)} \quad \nabla f(r) = f'(r) \nabla r$$

Soln.

$$\text{Given } \vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$$

$$\Rightarrow r = |\vec{r}| = \sqrt{x^2 + y^2 + z^2}$$

$$\Rightarrow r^2 = x^2 + y^2 + z^2 \rightarrow (1)$$

Difff. (1) w.r.t. to  $x, y, z$ ,

$$\begin{array}{l|l|l} \frac{\partial r}{\partial x} = \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} = \frac{\partial r}{\partial y} & \frac{\partial r}{\partial z} = \frac{\partial r}{\partial z} \\ \frac{\partial r}{\partial x} = \frac{x}{r} & \frac{\partial r}{\partial y} = \frac{y}{r} & \frac{\partial r}{\partial z} = \frac{z}{r} \end{array}$$

$$\begin{aligned} \text{i)} \quad \nabla r &= \vec{i} \frac{\partial r}{\partial x} + \vec{j} \frac{\partial r}{\partial y} + \vec{k} \frac{\partial r}{\partial z} \\ &= \vec{i} \left( \frac{x}{r} \right) + \vec{j} \left( \frac{y}{r} \right) + \vec{k} \left( \frac{z}{r} \right) \\ &= \frac{x\vec{i} + y\vec{j} + z\vec{k}}{r} \end{aligned}$$

$$\nabla r = \frac{\vec{r}}{r}$$

$$\begin{aligned} \text{ii)} \quad \nabla \left( \frac{1}{r} \right) &= \vec{i} \frac{\partial}{\partial x} \left( \frac{1}{r} \right) + \vec{j} \frac{\partial}{\partial y} \left( \frac{1}{r} \right) + \vec{k} \frac{\partial}{\partial z} \left( \frac{1}{r} \right) \\ &= \vec{i} \left[ -\frac{1}{r^2} \frac{\partial r}{\partial x} \right] + \vec{j} \left[ -\frac{1}{r^2} \frac{\partial r}{\partial y} \right] + \vec{k} \left[ -\frac{1}{r^2} \frac{\partial r}{\partial z} \right] \\ &= \vec{i} \left[ -\frac{1}{r^2} \times \frac{x}{r} \right] + \vec{j} \left[ -\frac{1}{r^2} \times \frac{y}{r} \right] + \vec{k} \left[ -\frac{1}{r^2} \times \frac{z}{r} \right] \\ &= -\frac{1}{r^3} [x\vec{i} + y\vec{j} + z\vec{k}] \end{aligned}$$



$\nabla \left( \frac{1}{r} \right) = - \frac{\vec{r}}{r^3}$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

$$\begin{aligned}
 \text{iii). } \nabla r^n &= \vec{i} \frac{\partial(r^n)}{\partial x} + \vec{j} \frac{\partial(r^n)}{\partial y} + \vec{k} \frac{\partial(r^n)}{\partial z} \\
 &= \vec{i} n r^{n-1} \frac{\partial r}{\partial x} + \vec{j} n r^{n-1} \frac{\partial r}{\partial y} + \vec{k} n r^{n-1} \frac{\partial r}{\partial z} \\
 &= n r^{n-1} \left[ \vec{i} \left( \frac{x}{r} \right) + \vec{j} \left( \frac{y}{r} \right) + \vec{k} \left( \frac{z}{r} \right) \right] \\
 &= \frac{n r^{n-1}}{r} \left[ x \vec{i} + y \vec{j} + z \vec{k} \right] \\
 &= \frac{n r^{n-1}}{r} \vec{r} \\
 \nabla r^n &= n r^{n-1} \vec{r}
 \end{aligned}$$

$$\begin{aligned}
 \text{iv). } \nabla f(r) &= \vec{i} \frac{\partial}{\partial x} f(r) + \vec{j} \frac{\partial}{\partial y} f(r) + \vec{k} \frac{\partial}{\partial z} f(r) \\
 &= \vec{i} f'(r) \frac{\partial r}{\partial x} + \vec{j} f'(r) \frac{\partial r}{\partial y} + \vec{k} f'(r) \frac{\partial r}{\partial z} \\
 &= f'(r) \left[ \vec{i} \left( \frac{x}{r} \right) + \vec{j} \left( \frac{y}{r} \right) + \vec{k} \left( \frac{z}{r} \right) \right] \\
 &= \frac{f'(r)}{r} \left[ x \vec{i} + y \vec{j} + z \vec{k} \right] \\
 &= f'(r) \times \frac{\vec{r}}{r}
 \end{aligned}$$

$$\nabla f(r) = f'(r) \nabla r \quad (\text{from iii})$$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

Surfaces :

$$\text{unit normal vector } \hat{n} = \frac{\nabla \phi}{|\nabla \phi|}$$

$$\text{normal derivative} = |\nabla \phi|$$

$$\text{directional derivative} = \nabla \phi \cdot \frac{\vec{a}}{|\vec{a}|}$$

Angle between the Surfaces :

$$\cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|}$$

If two surfaces are cut orthogonally, then  $\nabla \phi_1 \cdot \nabla \phi_2 = 0$

Q. Find the unit normal to the surface

$$x^2 + xy + z^2 = 4 \text{ at } (1, -1, 2).$$

Soln.

$$\text{Let } \phi = x^2 + xy + z^2 - 4$$

$$\text{unit normal vector } \hat{n} = \frac{\nabla \phi}{|\nabla \phi|}$$

Now

$$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$= \vec{i} \frac{\partial}{\partial x} (x^2 + xy + z^2 - 4) + \vec{j} \frac{\partial}{\partial y} (x^2 + xy + z^2 - 4) + \vec{k} \frac{\partial}{\partial z} (x^2 + xy + z^2 - 4)$$

$$= \vec{i} (2x + y) + \vec{j} (x) + \vec{k} (2z)$$

$$= \vec{i} (2(1) - 1) + \vec{j} (1) + \vec{k} (2(2))$$

$$\nabla \phi_{(1, -1, 2)}$$

$$= \vec{i} + \vec{j} + 4\vec{k}$$

$$\therefore \hat{n} = \frac{\vec{i} + \vec{j} + 4\vec{k}}{\sqrt{1+1+16}} = \frac{\vec{i} + \vec{j} + 4\vec{k}}{\sqrt{18}}$$

Q. Find the directional derivative of  $\phi = xyz$  at

Scalene with the direction of  $\vec{i} + \vec{j} + \vec{k}$





# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

## DERIVATIVES: Gradient of a scalar field, Directional Derivative

Soln.

$$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$\nabla \phi = \vec{i}(y) + \vec{j}(x) + \vec{k}(z)$$

$$\nabla \phi_{(1,1,1)} = \vec{i}(1) + \vec{j}(1) + \vec{k}(1)$$

$$= \vec{i} + \vec{j} + \vec{k}$$

Given  $\vec{a} = \vec{i} + \vec{j} + \vec{k}$

$$|\vec{a}| = \sqrt{1+1+1} = \sqrt{3}$$

$$DD = \nabla \phi \cdot \frac{\vec{a}}{|\vec{a}|}$$

$$= (\vec{i} + \vec{j} + \vec{k}) \cdot \frac{(\vec{i} + \vec{j} + \vec{k})}{\sqrt{3}}$$

$$= \frac{1+1+1}{\sqrt{3}} = \frac{3}{\sqrt{3}}$$

$$DD = \sqrt{3}$$

Q3. Find the directional derivative of  $\phi = x^2 + 2xy$  at  $(1, -1, 3)$  in the direction of  $\vec{i} + 2\vec{j} + 2\vec{k}$

Soln.

$$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$= \vec{i}(2x + 2y) + \vec{j}(2x) + \vec{k}(0)$$

$$\nabla \phi = (2x + 2y)\vec{i} + 2x\vec{j}$$

$$= (2(1) + 2(-1))\vec{i} + 2(-1)\vec{j}$$

$$\nabla \phi_{(1, -1, 3)} = -2\vec{j}$$

$$\vec{a} = \vec{i} + 2\vec{j} + 2\vec{k}$$

$$|\vec{a}| = \sqrt{1+4+4}$$

$$= \sqrt{9} = 3$$

$$DD = \nabla \phi \cdot \frac{\vec{a}}{|\vec{a}|} = -2\vec{j} \cdot \frac{\vec{i} + 2\vec{j} + 2\vec{k}}{3}$$

$$= -\frac{4}{3}$$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

## DERIVATIVES: Gradient of a scalar field, Directional Derivative

Q1. What is the greatest rate of increase of  $u = x^2 + yz^2$  at  $(1, -1, 3)$

Soln.

$$\nabla u = \vec{i} \frac{\partial u}{\partial x} + \vec{j} \frac{\partial u}{\partial y} + \vec{k} \frac{\partial u}{\partial z}$$

$$= \vec{i}(2x) + \vec{j}(z^2) + \vec{k}(2yz)$$

$$\nabla u = 2x\vec{i} + z^2\vec{j} + 2yz\vec{k}$$

$$\begin{aligned}\nabla u_{(1, -1, 3)} &= 2\vec{i} + 9\vec{j} + 2(-1)(3)\vec{k} \\ &= 2\vec{i} + 9\vec{j} - 6\vec{k}\end{aligned}$$

∴ The greatest rate increase in the direction of  $\vec{j}$ .

Q2. Find the angle b/w the normals to the surface  $xy = z^2$  at the points  $(1, 4, 2)$  &  $(-3, -3, 3)$

Soln.

$$\text{Given } xy = z^2$$

$$\phi = xy - z^2$$

$$\nabla \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$= \vec{i}(y) + \vec{j}(x) + \vec{k}(-2z)$$

$$= y\vec{i} + x\vec{j} - 2z\vec{k}$$

$$\nabla \phi_1 \text{ at } (1, 4, 2) = 4\vec{i} + \vec{j} - 4\vec{k}$$

$$|\nabla \phi_1| = \sqrt{16+1+16} = \sqrt{33}$$

$$\text{and } \nabla \phi_2 \text{ at } (-3, -3, 3) = -3\vec{i} - 3\vec{j} - 6\vec{k}$$

$$|\nabla \phi_2| = \sqrt{9+9+36} = \sqrt{54} = 3\sqrt{6}$$

$$\therefore \cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|}$$

$$= \frac{(4\vec{i} + \vec{j} - 4\vec{k}) \cdot (-3\vec{i} - 3\vec{j} - 6\vec{k})}{\sqrt{33} \cdot 3\sqrt{6}}$$

$$= \frac{-12 - 3 + 24}{3\sqrt{11 \times 3 \times 3 \times 2}} = \frac{9}{3 \times 3\sqrt{22}} = \frac{1}{\sqrt{22}}$$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

b). Find the angle between the surfaces

$$x^2 + y^2 - z^2 = 11 \text{ and } xy + yz - zx = 18 \text{ at } (6, 4, 3)$$

Soln.

$$\text{Let } \phi_1 = x^2 - y^2 - z^2 - 11$$

$$\begin{aligned} \nabla \phi_1 &= \vec{i} \frac{\partial \phi_1}{\partial x} + \vec{j} \frac{\partial \phi_1}{\partial y} + \vec{k} \frac{\partial \phi_1}{\partial z} \\ &= \vec{i}(2x) + \vec{j}(-2y) + \vec{k}(-2z) \end{aligned}$$

$$\nabla \phi_1(6, 4, 3) = 12\vec{i} - 8\vec{j} - 6\vec{k} \Rightarrow |\nabla \phi_1| = \sqrt{144 + 64 + 36} = \sqrt{244}$$

$$\text{and } \phi_2 = xy + yz - zx - 18 = \sqrt{244}$$

$$\begin{aligned} \nabla \phi_2 &= \vec{i} \frac{\partial \phi_2}{\partial x} + \vec{j} \frac{\partial \phi_2}{\partial y} + \vec{k} \frac{\partial \phi_2}{\partial z} \\ &= \vec{i}(y-z) + \vec{j}(x+z) + \vec{k}(y-x) \end{aligned}$$

$$\nabla \phi_2(6, 4, 3) = \vec{i} + 9\vec{j} - 2\vec{k} \Rightarrow |\nabla \phi_2| = \sqrt{1 + 81 + 4} = \sqrt{86}$$

$$\therefore \cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|}$$

$$= \frac{(12\vec{i} - 8\vec{j} - 6\vec{k}) \cdot (\vec{i} + 9\vec{j} - 2\vec{k})}{\sqrt{244} \sqrt{86}}$$

$$= \frac{12(1) - 8(9) - 6(-2)}{\sqrt{244} \sqrt{86}}$$

$$= \frac{-48}{2\sqrt{61} \sqrt{86}}$$

$$\cos \theta = \frac{-24}{\sqrt{5246}}$$

$$\theta = \cos^{-1} \left[ \frac{-24}{\sqrt{5246}} \right]$$

c). find a and b such that the surfaces  
 $ax^2 - byz = (a+2)x$  and  $4x^2y + z^3 = 4$  cut  
 orthogonally at  $(1, -1, 2)$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-I VECTOR CALCULUS

*sln.*

$$\text{Let } \phi_1 = ax^2 - byz - (a+2)x \rightarrow (1)$$

$$\nabla \phi_1 = \vec{i} \frac{\partial \phi_1}{\partial x} + \vec{j} \frac{\partial \phi_1}{\partial y} + \vec{k} \frac{\partial \phi_1}{\partial z}$$

$$\nabla \phi_1 = \vec{i} [2ax - (a+2)] + \vec{j} [-bz] + \vec{k} [-by]$$

$$\begin{aligned}\nabla \phi_1(1, -1, 2) &= \vec{i} [2a - a - 2] - 2b \vec{j} + b \vec{k} \\ &= (a-2) \vec{i} - 2b \vec{j} + b \vec{k}\end{aligned}$$

$$\text{and } \phi_2 = 4x^2y + z^3 - 4$$

$$\nabla \phi_2 = 8xy \vec{i} + 4x^2 \vec{j} + 3z^2 \vec{k}$$

$$\nabla \phi_2(1, -1, 2) = -8 \vec{i} + 4 \vec{j} + 12 \vec{k}$$

Given two surfaces are cut orthogonally,

$$\text{i.e., } \nabla \phi_1 \cdot \nabla \phi_2 = 0$$

$$[(a-2) \vec{i} - 2b \vec{j} + b \vec{k}] \cdot [-8 \vec{i} + 4 \vec{j} + 12 \vec{k}] = 0$$

$$-8(a-2) - 8b + 12b = 0$$

$$-8a + 16 - 8b + 12b = 0$$

$$-8a + b + 16 = 0$$

$$\text{i.e., } 8a - b - 16 = 0 \rightarrow (2)$$

Since  $(1, -1, 2)$  lies on the surface using (1).

$$\phi_1(x, y, z) = 0$$

$$a(1)^2 - b(-1)(2) = (a+2)(1)$$

$$a + 2b - a - 2 = 0$$

$$2b = 2 \Rightarrow \boxed{b=1}$$

$$(2) \Rightarrow 8a - 1 - 4 = 0 \Rightarrow 8a = 5 \Rightarrow \boxed{a = 5/8}$$



Scanned with  
CamScanner