



2]. Solve $(3x+2)^2 \frac{d^2y}{dx^2} + 3(3x+2) \frac{dy}{dx} - 36y = 3x^2 + 4x + 1$

Soln.

Given $[(3x+2)^2 D^2 + 3(3x+2) D - 36] y = 3x^2 + 4x + 1$

Take $3x+2 = e^x \Rightarrow 3x = e^x - 2 \Rightarrow x = \frac{e^x - 2}{3} \rightarrow (1)$

$\log(3x+2) = x$

$(3x+2) D = 3D'$

$(3x+2)^2 D^2 = 9D'(D'-1)$

(1) $\Rightarrow [9D'(D'-1) + 3(3D') - 36] y = 3 \left[\frac{e^x - 2}{3} \right]^2 + 4 \left[\frac{e^x - 2}{3} \right] + 1$

$[9D'^2 - 9D' + 9D' - 36] y = \frac{3}{9} [e^{2x} + 4 - 4e^x] + \frac{4}{3} e^x - \frac{8}{3} + 1$

$[9D'^2 - 36] y = \frac{e^{2x}}{3} + \frac{4}{3} - \frac{4e^x}{3} + \frac{4}{3} e^x - \frac{8}{3} + 1$

$= \frac{e^{2x}}{3} - \frac{1}{3}$

$9(D'^2 - 4) y = \frac{1}{3} [e^{2x} - 1]$

$(D'^2 - 4) y = \frac{1}{27} [e^{2x} - 1]$

CF:

$m^2 - 4 = 0 \Rightarrow m^2 = 4$
 $m = \pm 2$

CF = $Ae^{2x} + Be^{-2x}$

PI = $\frac{1}{D^2 - 4} \left[\frac{1}{27} (e^{2x} - 1) \right]$

$= \frac{1}{27} \left[\frac{1}{D^2 - 4} e^{2x} - \frac{1}{D^2 - 4} e^{0x} \right]$

$= \frac{1}{27} \left[x \frac{1}{2D'} e^{2x} - \frac{1}{(-4)} e^{0x} \right] \quad \begin{matrix} 1 \rightarrow D' \rightarrow 2 \\ 2 \rightarrow D' \rightarrow 0 \end{matrix}$

$= \frac{1}{27} \left[\frac{x}{4} e^{2x} + \frac{1}{4} \right]$

$= \frac{1}{108} [xe^{2x} + 1]$

The soln. is, $y = CF + PI = Ae^{2x} + Be^{-2x} + \frac{1}{108} [xe^{2x} + 1]$



Scanned with CamScanner

$y = A(3x+2)^2 + B(3x+2)^{-2} + \frac{1}{108} [\log(3x+2) \{3x+2\}^2 + 1]$