



Unit - II

Vector calculus

Gradient:

Let $\phi(x, y, z)$ be a scalar point function and is continuously differentiable. Then the vector

$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$ is called the gradient of the scalar fn. ϕ .

i.e., $\text{grad } \phi = \nabla\phi$

Problems

1] Find $\nabla\phi$ where $\phi = x^2 + y^2 + z^2$

Soln.

$$\text{Grad. } \phi \text{ (or) } \nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$= \vec{i} \frac{\partial(x^2 + y^2 + z^2)}{\partial x} + \vec{j} \frac{\partial(x^2 + y^2 + z^2)}{\partial y} + \vec{k} \frac{\partial(x^2 + y^2 + z^2)}{\partial z}$$

$$= \vec{i} (2x) + \vec{j} (2y) + \vec{k} (2z)$$

$$\nabla\phi = 2x\vec{i} + 2y\vec{j} + 2z\vec{k}$$

2]. Find $\nabla\phi$ where $\phi = 3x^2y - y^3z^2$ at (1, 1, 1)

Soln.

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$= \vec{i} \frac{\partial(3x^2y - y^3z^2)}{\partial x} + \vec{j} \frac{\partial(3x^2y - y^3z^2)}{\partial y} + \vec{k} \frac{\partial(3x^2y - y^3z^2)}{\partial z}$$

$$= \vec{i} [6xy - 0] + \vec{j} [3x^2 - 3y^3z^2] + \vec{k} [0 - 2y^3z]$$

$$\nabla\phi = 6xy\vec{i} + (3x^2 - 3y^3z^2)\vec{j} + 2y^3z\vec{k}$$

$$\nabla\phi_{(1,1,1)} = 6(1)(1)\vec{i} + (3 - 3)\vec{j} - 2(1)(1)\vec{k}$$

$$= 6\vec{i} + 0\vec{j} - 2\vec{k}$$



$$= 6\vec{i} - 2\vec{k}$$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

3]. Find the maximum directional derivative of $\phi = xyz^2$ at $(1, 0, 3)$

Soln.

$$\begin{aligned}\nabla\phi &= \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z} \\ &= \vec{i} \frac{\partial}{\partial x} (xyz^2) + \vec{j} \frac{\partial}{\partial y} (xyz^2) + \vec{k} \frac{\partial}{\partial z} (xyz^2)\end{aligned}$$

$$\nabla\phi = \vec{i}(yz^2) + \vec{j}(xz^2) + \vec{k}(2xyz)$$

$$\begin{aligned}\nabla\phi_{(1,0,3)} &= \vec{i}(0) + \vec{j}(9) + \vec{k}(0) \\ &= 9\vec{j} \quad \text{maximum D.D} = \sqrt{9} = 3\end{aligned}$$

4]. Find $\nabla\phi$ where $\phi = xyz$ at $(1, 2, 3)$

Soln.

$$\begin{aligned}\nabla\phi &= \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z} \\ &= \vec{i} \frac{\partial}{\partial x} (xyz) + \vec{j} \frac{\partial}{\partial y} (xyz) + \vec{k} \frac{\partial}{\partial z} (xyz)\end{aligned}$$

$$\nabla\phi = \vec{i}(yz) + \vec{j}(xz) + \vec{k}(xy)$$

$$\begin{aligned}\nabla\phi_{(1,2,3)} &= \vec{i}(2)(3) + \vec{j}(1)(3) + \vec{k}(1)(2) \\ &= 6\vec{i} + 3\vec{j} + 2\vec{k}\end{aligned}$$

5]. If $\nabla\phi = yz\vec{i} + zx\vec{j} + xy\vec{k}$, find ϕ .

Soln.

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$\nabla\phi = \vec{i}(yz) + \vec{j}(xz) + \vec{k}(xy)$$

Equating w.r. to $\vec{i}, \vec{j}, \vec{k}$

$$\frac{\partial\phi}{\partial x} = yz \quad \left| \quad \frac{\partial\phi}{\partial y} = zx \quad \left| \quad \frac{\partial\phi}{\partial z} = xy \right. \right.$$

Integrate w.r. to x w.r. to y w.r. to z

$$\phi = xyz + f(y, z) \quad \left| \quad \phi = xyz + f(x, z) \quad \left| \quad \phi = xyz + f(x, y) \right. \right.$$

In general,

$$\phi = xyz + C$$



Scanned with CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

6J. If $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$, such that $|\vec{r}| = r$,

prove that

i). $\nabla r = \frac{\vec{r}}{r} = \hat{r}$

iii). $\nabla r^n = n r^{n-2} \vec{r}$

ii). $\nabla(\frac{1}{r}) = \frac{-\vec{r}}{r^3} = \frac{-\hat{r}}{r^2}$

iv). $\nabla f(r) = f'(r) \nabla r$

Soln.

Given $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$

$\Rightarrow r = |\vec{r}| = \sqrt{x^2 + y^2 + z^2}$

$\Rightarrow r^2 = x^2 + y^2 + z^2 \rightarrow (1)$

Diff. (1) w.r. to x, y, z ,

$$2r \frac{\partial r}{\partial x} = 2x \quad \left| \quad 2r \frac{\partial r}{\partial y} = 2y \quad \right| \quad 2r \frac{\partial r}{\partial z} = 2z$$

$$\frac{\partial r}{\partial x} = \frac{x}{r} \quad \left| \quad \frac{\partial r}{\partial y} = \frac{y}{r} \quad \right| \quad \frac{\partial r}{\partial z} = \frac{z}{r}$$

i). $\nabla r = \vec{i} \frac{\partial r}{\partial x} + \vec{j} \frac{\partial r}{\partial y} + \vec{k} \frac{\partial r}{\partial z}$

$$= \vec{i} \left(\frac{x}{r}\right) + \vec{j} \left(\frac{y}{r}\right) + \vec{k} \left(\frac{z}{r}\right)$$

$$= \frac{x\vec{i} + y\vec{j} + z\vec{k}}{r}$$

$\nabla r = \frac{\vec{r}}{r}$

ii). $\nabla(\frac{1}{r}) = \vec{i} \frac{\partial}{\partial x}(\frac{1}{r}) + \vec{j} \frac{\partial}{\partial y}(\frac{1}{r}) + \vec{k} \frac{\partial}{\partial z}(\frac{1}{r})$

$$= \vec{i} \left[-\frac{1}{r^2} \frac{\partial r}{\partial x}\right] + \vec{j} \left[-\frac{1}{r^2} \frac{\partial r}{\partial y}\right] + \vec{k} \left[-\frac{1}{r^2} \frac{\partial r}{\partial z}\right]$$

$$= \vec{i} \left[-\frac{1}{r^2} \times \frac{x}{r}\right] + \vec{j} \left[-\frac{1}{r^2} \times \frac{y}{r}\right] + \vec{k} \left[-\frac{1}{r^2} \times \frac{z}{r}\right]$$

$$= -\frac{1}{r^3} [x\vec{i} + y\vec{j} + z\vec{k}]$$

$\nabla(\frac{1}{r}) = \frac{-\vec{r}}{r^3}$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

$$\begin{aligned} \text{iii). } \nabla r^n &= \vec{i} \frac{\partial (r^n)}{\partial x} + \vec{j} \frac{\partial (r^n)}{\partial y} + \vec{k} \frac{\partial (r^n)}{\partial z} \\ &= \vec{i} n r^{n-1} \frac{\partial r}{\partial x} + \vec{j} n r^{n-1} \frac{\partial r}{\partial y} + \vec{k} n r^{n-1} \frac{\partial r}{\partial z} \\ &= n r^{n-1} \left[\vec{i} \frac{\partial r}{\partial x} + \vec{j} \frac{\partial r}{\partial y} + \vec{k} \frac{\partial r}{\partial z} \right] \\ &= n r^{n-1} \left[\vec{i} \left(\frac{x}{r} \right) + \vec{j} \left(\frac{y}{r} \right) + \vec{k} \left(\frac{z}{r} \right) \right] \\ &= \frac{n r^{n-1}}{r} [x \vec{i} + y \vec{j} + z \vec{k}] \\ &= \frac{n r^{n-1}}{r} \vec{r} \\ \nabla r^n &= n r^{n-2} \vec{r} \end{aligned}$$

$$\begin{aligned} \text{iv). } \nabla f(r) &= \vec{i} \frac{\partial}{\partial x} f(r) + \vec{j} \frac{\partial}{\partial y} f(r) + \vec{k} \frac{\partial}{\partial z} f(r) \\ &= \vec{i} f'(r) \frac{\partial r}{\partial x} + \vec{j} f'(r) \frac{\partial r}{\partial y} + \vec{k} f'(r) \frac{\partial r}{\partial z} \\ &= f'(r) \left[\vec{i} \left(\frac{x}{r} \right) + \vec{j} \left(\frac{y}{r} \right) + \vec{k} \left(\frac{z}{r} \right) \right] \\ &= \frac{f'(r)}{r} [x \vec{i} + y \vec{j} + z \vec{k}] \\ &= f'(r) \times \frac{\vec{r}}{r} \end{aligned}$$

$$\nabla f(r) = f'(r) \nabla r \quad (\text{from (i)})$$



Scanned with
CamScanner



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

Surfaces:

$$\text{Unit normal vector } \hat{n} = \frac{\nabla\phi}{|\nabla\phi|}$$

$$\text{Normal derivative} = |\nabla\phi|$$

$$\text{Directional derivative} = \nabla\phi \cdot \frac{\vec{a}}{|\vec{a}|}$$

Angle between the Surfaces:

$$\cos \theta = \frac{\nabla\phi_1 \cdot \nabla\phi_2}{|\nabla\phi_1| |\nabla\phi_2|}$$

If two surfaces are cut orthogonally, then $\nabla\phi_1 \cdot \nabla\phi_2 = 0$

Q. Find the unit normal to the surface

$$x^2 + xy + z^2 = 4 \text{ at } (1, -1, 2).$$

Soln.

$$\text{Let } \phi = x^2 + xy + z^2 - 4$$

$$\text{unit normal vector } \hat{n} = \frac{\nabla\phi}{|\nabla\phi|}$$

$$\text{Now } \nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$= \vec{i} \frac{\partial}{\partial x} (x^2 + xy + z^2 - 4) + \vec{j} \frac{\partial}{\partial y} (x^2 + xy + z^2 - 4) + \vec{k} \frac{\partial}{\partial z} (x^2 + xy + z^2 - 4)$$

$$= \vec{i} (2x + y) + \vec{j} (x) + \vec{k} (2z)$$

$$\nabla\phi_{(1, -1, 2)} = \vec{i} (2(1) - 1) + \vec{j} (1) + \vec{k} (2(2))$$

$$= \vec{i} + \vec{j} + 4\vec{k}$$

$$\therefore \hat{n} = \frac{\vec{i} + \vec{j} + 4\vec{k}}{\sqrt{1+1+16}} = \frac{\vec{i} + \vec{j} + 4\vec{k}}{\sqrt{18}}$$

2]. Find the directional derivative of $\phi = xyz$ at

in the direction of $\vec{i} + \vec{j} + \vec{k}$



Scanned with CamScanner



Soln.

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$\nabla\phi = \vec{i}(yx) + \vec{j}(xz) + \vec{k}(xy)$$

$$\nabla\phi_{(1,1,1)} = \vec{i}(1)(1) + \vec{j}(1)(1) + \vec{k}(1)(1)$$

$$= \vec{i} + \vec{j} + \vec{k}$$

Given $\vec{a} = \vec{i} + \vec{j} + \vec{k}$

$$|\vec{a}| = \sqrt{1+1+1} = \sqrt{3}$$

$$DD = \nabla\phi \cdot \frac{\vec{a}}{|\vec{a}|}$$

$$= (\vec{i} + \vec{j} + \vec{k}) \cdot \frac{(\vec{i} + \vec{j} + \vec{k})}{\sqrt{3}}$$

$$= \frac{1+1+1}{\sqrt{3}} = \frac{3}{\sqrt{3}}$$

$$DD = \sqrt{3}$$

3]. Find the directional derivative of $\phi = x^2 + 2xy$ at $(1, -1, 3)$ in the direction of $\vec{i} + 2\vec{j} + 2\vec{k}$

Soln.

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$= \vec{i}(2x+2y) + \vec{j}(2x) + \vec{k}(0)$$

$$\nabla\phi = (2x+2y)\vec{i} + 2x\vec{j}$$

$$\nabla\phi_{(1,-1,3)} = (2(1)+2(-1))\vec{i} + 2(1)\vec{j}$$

$$= -2\vec{j}$$

$$\vec{a} = \vec{i} + 2\vec{j} + 2\vec{k}$$

$$|\vec{a}| = \sqrt{1+4+4}$$

$$= \sqrt{9} = 3$$

$$DD = \nabla\phi \cdot \frac{\vec{a}}{|\vec{a}|} = -2\vec{j} \cdot \frac{\vec{i} + 2\vec{j} + 2\vec{k}}{3}$$

$$= -\frac{4}{3}$$





SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

4]. What is the greatest rate of increase of $u = x^2 + yz^2$ at $(1, -1, 3)$

Soln.

$$\begin{aligned}\nabla u &= \vec{i} \frac{\partial u}{\partial x} + \vec{j} \frac{\partial u}{\partial y} + \vec{k} \frac{\partial u}{\partial z} \\ &= \vec{i} (2x) + \vec{j} (z^2) + \vec{k} (2yz)\end{aligned}$$

$$\nabla u = 2x \vec{i} + z^2 \vec{j} + 2yz \vec{k}$$

$$\begin{aligned}\nabla u_{(1, -1, 3)} &= 2\vec{i} + 9\vec{j} + 2(-1)(3)\vec{k} \\ &= 2\vec{i} + 9\vec{j} - 6\vec{k}\end{aligned}$$

\therefore The greatest rate increase in the direction of \vec{j} .

5]. Find the angle b/w the normals to the surface $xy = z^2$ at the points $(1, 4, 2)$ & $(-3, -3, 3)$

Soln.

Given $xy = z^2$

$$\phi = xy - z^2$$

$$\begin{aligned}\nabla \phi &= \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z} \\ &= \vec{i} (y) + \vec{j} (x) + \vec{k} (-2z) \\ &= y \vec{i} + x \vec{j} - 2z \vec{k}\end{aligned}$$

$$\nabla \phi_1 (1, 4, 2) = 4\vec{i} + \vec{j} - 4\vec{k}$$

$$|\nabla \phi_1| = \sqrt{16 + 1 + 16} = \sqrt{33}$$

$$\text{and } \nabla \phi_2 (-3, -3, 3) = -3\vec{i} - 3\vec{j} - 6\vec{k}$$

$$|\nabla \phi_2| = \sqrt{9 + 9 + 36} = \sqrt{54} = 3\sqrt{6}$$

$$\therefore \cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|}$$

$$= \frac{(4\vec{i} + \vec{j} - 4\vec{k}) \cdot (-3\vec{i} - 3\vec{j} - 6\vec{k})}{\sqrt{33} \cdot 3\sqrt{6}}$$

$$= \frac{-12 - 3 + 24}{3\sqrt{11 \times 3 \times 3 \times 2}} = \frac{9}{3 \times 3\sqrt{22}} = \frac{1}{\sqrt{22}}$$



Scanned with CamScanner $\rightarrow \theta = \cos^{-1} \left(\frac{1}{\sqrt{22}} \right)$



SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-I VECTOR CALCULUS

DERIVATIVES: Gradient of a scalar field, Directional Derivative

6]. Find the angle between the surfaces

$$x^2 - y^2 - z^2 = 11 \text{ and } xy + yz - zx = 18 \text{ at } (6, 4, 3)$$

Soln.

$$\text{Let } \phi_1 = x^2 - y^2 - z^2 - 11$$

$$\begin{aligned} \nabla \phi_1 &= \vec{i} \frac{\partial \phi_1}{\partial x} + \vec{j} \frac{\partial \phi_1}{\partial y} + \vec{k} \frac{\partial \phi_1}{\partial z} \\ &= \vec{i}(2x) + \vec{j}(-2y) + \vec{k}(-2z) \end{aligned}$$

$$\nabla \phi_1(6, 4, 3) = 12\vec{i} - 8\vec{j} - 6\vec{k} \Rightarrow |\nabla \phi_1| = \sqrt{144 + 64 + 36} = \sqrt{244}$$

and $\phi_2 = xy + yz - zx - 18$

$$\begin{aligned} \nabla \phi_2 &= \vec{i} \frac{\partial \phi_2}{\partial x} + \vec{j} \frac{\partial \phi_2}{\partial y} + \vec{k} \frac{\partial \phi_2}{\partial z} \\ &= \vec{i}(y - z) + \vec{j}(x + z) + \vec{k}(y - x) \end{aligned}$$

$$\nabla \phi_2(6, 4, 3) = \vec{i} + 9\vec{j} - 3\vec{k} \Rightarrow |\nabla \phi_2| = \sqrt{1 + 81 + 9} = \sqrt{91}$$

$$\begin{aligned} \therefore \cos \theta &= \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|} \\ &= \frac{(12\vec{i} - 8\vec{j} - 6\vec{k}) \cdot (\vec{i} + 9\vec{j} - 3\vec{k})}{\sqrt{244} \sqrt{91}} \\ &= \frac{12(1) - 8(9) - 6(-2)}{\sqrt{244} \sqrt{91}} \\ &= \frac{-48}{2\sqrt{61} \sqrt{91}} \end{aligned}$$

$$\cos \theta = \frac{-24}{\sqrt{5246}}$$

$$\theta = \cos^{-1} \left[\frac{-24}{\sqrt{5246}} \right]$$

7]. Find a and b, such that the surfaces $ax^2 - byz = (a+2)x$ and $4x^2y + z^3 = 4$ cut orthogonally at $(1, -1, 2)$



Scanned with CamScanner



Soln.

$$\text{Let } \phi_1 = ax^2 - byz - (a+z)x \rightarrow (1)$$

$$\nabla \phi_1 = \vec{i} \frac{\partial \phi_1}{\partial x} + \vec{j} \frac{\partial \phi_1}{\partial y} + \vec{k} \frac{\partial \phi_1}{\partial z}$$

$$\nabla \phi_1 = \vec{i} [2ax - (a+z)] + \vec{j} [-bz] + \vec{k} [-by]$$

$$\nabla \phi_1 (1, -1, 2) = \vec{i} [2a - a - 2] - 2b\vec{j} + b\vec{k}$$

$$= (a-2)\vec{i} - 2b\vec{j} + b\vec{k}$$

$$\text{and } \phi_2 = 4x^2y + z^3 - 4$$

$$\nabla \phi_2 = 8xy\vec{i} + 4x^2\vec{j} + 3z^2\vec{k}$$

$$\nabla \phi_2 (1, -1, 2) = -8\vec{i} + 4\vec{j} + 12\vec{k}$$

Given two surfaces are cut orthogonally,

$$\text{i.e., } \nabla \phi_1 \cdot \nabla \phi_2 = 0$$

$$[(a-2)\vec{i} - 2b\vec{j} + b\vec{k}] \cdot [-8\vec{i} + 4\vec{j} + 12\vec{k}] = 0$$

$$-8(a-2) - 8b + 12b = 0$$

$$-8a + 16 - 8b + 12b = 0$$

$$-2a + b + 4 = 0$$

$$\text{i.e., } 2a - b - 4 = 0 \rightarrow (2)$$

Since (1, -1, 2) lies on the surface using (1).

$$\phi_1(x, y, z) = 0$$

$$a(1)^2 - b(-1)(2) = (a+2)(1)$$

$$a + 2b - a - 2 = 0$$

$$2b = 2 \Rightarrow \boxed{b=1}$$

$$(2) \Rightarrow 2a - 1 - 4 = 0 \Rightarrow 2a = 5 \Rightarrow \boxed{a=5/2}$$

