



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS



## PART A

1. If  $\phi = x^2 + y^2 + z^2$ , find  $\nabla\phi$  at  $(1,1,-1)$

Given,  $\phi = x^2 + y^2 + z^2$  ----- (i)

There fore  $\nabla\phi = \vec{i}\frac{\partial\phi}{\partial x} + \vec{j}\frac{\partial\phi}{\partial y} + \vec{k}\frac{\partial\phi}{\partial z}$  ----- (ii)

From (i),  $\frac{\partial\phi}{\partial x} = 2x$  ;  $\frac{\partial\phi}{\partial y} = 2y$  ;  $\frac{\partial\phi}{\partial z} = 2z$  -----(iii)

Sub (iii) in (i) , we get

$$\nabla\phi = \vec{i}(2x) + \vec{j}(2y) + \vec{k}(2z)$$

There fore,  $(\nabla\phi)_{at(1,1,-1)} = 2\vec{i} + 2\vec{j} - 2\vec{k}$

2. Find grad  $r^n$ , where  $r = |\vec{r}|$  and  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$

Given,  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$

$$|\vec{r}| = r = \sqrt{x^2 + y^2 + z^2}$$

$$r^2 = x^2 + y^2 + z^2$$
 ----- (i)

Diff (i) partially w.r.t 'x'

$$2r \frac{\partial r}{\partial x} = 2x$$

$$\Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}$$

$$\frac{\partial r}{\partial y} = \frac{y}{r} \text{ and } \frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\therefore \text{grad}r^n = \nabla r^n$$

$$= \sum \vec{i} \frac{\partial}{\partial x} (r^n)$$

$$= \sum \vec{i} \frac{\partial}{\partial x} (r^n) \cdot \frac{\partial r}{\partial x}$$

$$= \vec{i} \cdot n \cdot r^{n-1} \cdot \frac{x}{r}$$

$$= \vec{i} \cdot n \cdot r^{n-2} \cdot x$$

$$= n \cdot r^{n-2} [x\vec{i} + y\vec{j} + z\vec{k}]$$

$$= n \cdot r^{n-2} \vec{r}$$



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

**3. Find the unit vector normal to the surface  $x^2 + y^2 - z = 10$  at  $(1,1,1)$ .**

**Given**  $\phi = x^2 + y^2 - z = 10$

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

$$= 2x\vec{i} + 2y\vec{j} - \vec{k}$$

$$\therefore (\nabla\phi)_{\text{at}(1,1,1)} = 2\vec{i} + 2\vec{j} - \vec{k}$$

$$|\nabla\phi| = \sqrt{4+4+1} = \sqrt{9} = 3$$

**Unit normal vector**  $\hat{n} = \frac{\nabla\phi}{|\nabla\phi|} = \frac{2\vec{i} + 2\vec{j} - \vec{k}}{3}$

**4. Find the directional derivative of  $\phi = xy + yz + xz$  at the point  $(1,2,3)$  in the direction  $3\vec{i} + 4\vec{j} + 5\vec{k}$ .**

**Given,**  $\phi = xy + yz + xz$  -----(i)

**Let**  $\vec{n} = 3\vec{i} + 4\vec{j} + 5\vec{k}$  ----- (ii)

**Directional derivative** =  $(\nabla\phi) \cdot \hat{n}$  ----- (A)

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

**From (i),**  $\frac{\partial\phi}{\partial x} = y + z$  ;  $\frac{\partial\phi}{\partial y} = x + z$  ;  $\frac{\partial\phi}{\partial z} = y + x$

$$\therefore \nabla\phi = (y + z)\vec{i} + (x + z)\vec{j} + (y + x)\vec{k}$$

$$\therefore (\nabla\phi)_{\text{at}(1,2,3)} = 5\vec{i} + 4\vec{j} + 3\vec{k}$$

**From (ii), we have**

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{3\vec{i} + 4\vec{j} + 5\vec{k}}{\sqrt{50}} \text{ ----- (iv)}$$

**Sub (iii) and (iv) in (A), we get**

$$\text{Directional derivative} = (\nabla\phi) \cdot \hat{n} = (5\vec{i} + 4\vec{j} + 3\vec{k}) \cdot \frac{3\vec{i} + 4\vec{j} + 5\vec{k}}{\sqrt{50}}$$

$$= \frac{15 + 16 + 15}{\sqrt{25 \times 2}} = \frac{46}{5\sqrt{2}}$$



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---

5. In what direction from the point (1,-1,-2) is the directional derivative of  $\phi = x^3y^3z^3$  a maximum? What is the magnitude of this maximum?

Given,  $\phi = x^3y^3z^3$  -----(i)

$$\nabla\phi = \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z}$$

From (i),  $\frac{\partial\phi}{\partial x} = 3x^2y^3z^3$  ;  $\frac{\partial\phi}{\partial y} = 3x^3y^2z^3$  ;  $\frac{\partial\phi}{\partial z} = 3x^3y^3z^2$

$$\therefore \nabla\phi = 3x^2y^3z^3\vec{i} + 3x^3y^2z^3\vec{j} + 3x^3y^3z^2\vec{k}$$

$$\therefore (\nabla\phi)_{at(1,2,3)} = 24\vec{i} - 24\vec{j} - 12\vec{k}$$

There fore the directional derivative is maximum in the direction  $24\vec{i} - 24\vec{j} - 12\vec{k}$ .

Magnitude of this maximum is  $|\nabla\phi|$

$$\begin{aligned} &= \sqrt{(24)^2 + (-24)^2 + (-12)^2} \\ &= \sqrt{1296} = 36 \end{aligned}$$

6. Find the angle between the normal to the surface  $xy = z^2$  at the points (1,4,2) and (-3,-3,3).

Let  $\phi = xy - z^2$  -----(i)

$$\therefore \nabla\phi = y\vec{i} + x\vec{j} - 2z\vec{k}$$

Normal to the surface is  $\nabla_1\phi$  and  $\nabla_2\phi$

$$\therefore \nabla_1\phi = (\nabla\phi)_{at(1,4,2)} = 4\vec{i} + \vec{j} - 4\vec{k}$$

$$\nabla_2\phi = (\nabla\phi)_{at(-3,-3,3)} = -3\vec{i} - 3\vec{j} - 6\vec{k}$$

$$\therefore |\nabla_1\phi| = \sqrt{33}; |\nabla_2\phi| = \sqrt{54}$$

There fore angle between the normal to the surface is,

$$\begin{aligned} \cos\theta &= \frac{(\nabla_1\phi)(\nabla_2\phi)}{|\nabla_1\phi||\nabla_2\phi|} = \frac{(4\vec{i} + \vec{j} - 4\vec{k}) \cdot (-3\vec{i} - 3\vec{j} - 6\vec{k})}{\sqrt{33}\sqrt{54}} \\ &= \frac{9}{\sqrt{1782}} = \frac{9}{9\sqrt{22}} = \frac{1}{\sqrt{22}} \\ \therefore \theta &= \cos^{-1}\left[\frac{1}{\sqrt{22}}\right] \end{aligned}$$

7. If  $\phi$  is a scalar point function, then prove that  $\text{curl}(\text{grad}\phi) = 0$ .



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

$$\text{grad } \phi = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

$$\text{curl grad } \phi = \nabla \times \left[ \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z} \right]$$

$$= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial \phi}{\partial x} & \frac{\partial \phi}{\partial y} & \frac{\partial \phi}{\partial z} \end{vmatrix}$$

$$= \vec{i} \left[ \frac{\partial^2 \phi}{\partial y \partial z} - \frac{\partial^2 \phi}{\partial y \partial z} \right] - \vec{j} \left[ \frac{\partial^2 \phi}{\partial x \partial z} - \frac{\partial^2 \phi}{\partial x \partial z} \right] + \vec{k} \left[ \frac{\partial^2 \phi}{\partial x \partial y} - \frac{\partial^2 \phi}{\partial x \partial y} \right]$$

$$= \mathbf{0}$$

**8. If  $\vec{A}$  is a constant vector, prove that  $\text{div } \vec{A} = 0$ .**

Let  $\vec{A} = A_1 \vec{i} + A_2 \vec{j} + A_3 \vec{k}$

Where  $A_1, A_2, A_3$  are constants

$$\therefore \text{div } \vec{A} = \nabla \cdot \vec{A}$$

$$= \left( \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \cdot (A_1 \vec{i} + A_2 \vec{j} + A_3 \vec{k})$$

$$= \frac{\partial A_1}{\partial x} + \frac{\partial A_2}{\partial y} + \frac{\partial A_3}{\partial z} = \mathbf{0+0+0}$$

$$\text{div } \vec{A} = \mathbf{0}$$

**9. If  $\vec{A}$  is a constant vector, prove that  $\text{curl } \vec{A} = 0$ .**

Let  $\vec{A} = A_1 \vec{i} + A_2 \vec{j} + A_3 \vec{k}$

Where  $A_1, A_2, A_3$  are constants

$$\text{curl } \vec{A} = \nabla \times \vec{A} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_1 & A_2 & A_3 \end{vmatrix}$$

$$= \vec{i} \left[ \frac{\partial A_3}{\partial y} - \frac{\partial A_2}{\partial z} \right] - \vec{j} \left[ \frac{\partial A_3}{\partial x} - \frac{\partial A_1}{\partial z} \right] + \vec{k} \left[ \frac{\partial A_2}{\partial x} - \frac{\partial A_1}{\partial y} \right]$$

$$= \vec{i}(0-0) - \vec{j}(0-0) + \vec{k}(0-0)$$

$$\text{curl } \vec{A} = \mathbf{0}$$

**10. Determine  $f(\mathbf{r})$  so that the vector  $f(\mathbf{r}) \vec{r}$  is solenoidal.**



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

Since  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$

$$\mathbf{f}(\mathbf{r}) = x\mathbf{f}'(\mathbf{r})\vec{i} + y\mathbf{f}'(\mathbf{r})\vec{j} + z\mathbf{f}'(\mathbf{r})\vec{k}$$

$$\text{div} [\mathbf{f}(\mathbf{r})] = \frac{\partial}{\partial x}[x\mathbf{f}'(\mathbf{r})] + \frac{\partial}{\partial y}[y\mathbf{f}'(\mathbf{r})] + \frac{\partial}{\partial z}[z\mathbf{f}'(\mathbf{r})]$$

$$= \mathbf{f}'(\mathbf{r}) + x\mathbf{f}''(\mathbf{r})\frac{\partial \mathbf{r}}{\partial x} + y\mathbf{f}''(\mathbf{r})\frac{\partial \mathbf{r}}{\partial y} + z\mathbf{f}''(\mathbf{r})\frac{\partial \mathbf{r}}{\partial z}$$

$$= 3\mathbf{f}'(\mathbf{r}) + \mathbf{f}''(\mathbf{r}) \left[ x\frac{\partial \mathbf{r}}{\partial x} + y\frac{\partial \mathbf{r}}{\partial y} + z\frac{\partial \mathbf{r}}{\partial z} \right]$$

$$= 3\mathbf{f}'(\mathbf{r}) + \mathbf{f}''(\mathbf{r}) \left[ x\frac{x}{r} + y\frac{y}{r} + z\frac{z}{r} \right]$$

$$= 3\mathbf{f}'(\mathbf{r}) + \frac{\mathbf{f}''(\mathbf{r})}{r} [x^2 + y^2 + z^2]$$

$$= 3\mathbf{f}'(\mathbf{r}) + r\mathbf{f}''(\mathbf{r})$$

Since  $\mathbf{f}(\mathbf{r})\vec{r}$  is solenoidal,  $\text{div}[\mathbf{f}(\mathbf{r})\vec{r}] = 0$

$$\text{ie., } 3\mathbf{f}'(\mathbf{r}) + r\mathbf{f}''(\mathbf{r}) = 0$$

$$\frac{\mathbf{f}''(\mathbf{r})}{\mathbf{f}'(\mathbf{r})} = \frac{-3}{r}$$

Integrating w.r.t  $r$ , we get

$$\log \mathbf{f}'(\mathbf{r}) = -3\log r + \log c$$

$$\log \mathbf{f}'(\mathbf{r}) = \log cr^{-3}$$

$$\mathbf{f}'(\mathbf{r}) = cr^{-3}$$

$$\mathbf{f}(\mathbf{r}) = \frac{c}{r^3}$$

**11. Find the value of 'a' so that the vector,  $\vec{F} = (x + 3y)\vec{i} + (y - 2z)\vec{j} + (x + az)\vec{k}$  is Solenoidal.**

Given  $\vec{F}$  is solenoidal.

$$\text{div } \vec{F} = 0$$

$$\text{ie., } \nabla \cdot \vec{F} = 0$$

$$\text{ie., } \left( \vec{i}\frac{\partial}{\partial x} + \vec{j}\frac{\partial}{\partial y} + \vec{k}\frac{\partial}{\partial z} \right) \cdot [(x + 3y)\vec{i} + (y - 2z)\vec{j} + (x + az)\vec{k}] = 0$$

$$\text{ie., } \frac{\partial}{\partial x}(x + 3y) + \frac{\partial}{\partial y}(y - 2z) + \frac{\partial}{\partial z}(x + az) = 0 \Rightarrow 1 + 1 + a = 0 \Rightarrow a = -2$$

**12. Show that the vector  $2xy\vec{i} + (x^2 + 2yz)\vec{j} + (y^2 + 1)\vec{k}$  is irrotational.**

$$\text{Let } \vec{F} = 2xy\vec{i} + (x^2 + 2yz)\vec{j} + (y^2 + 1)\vec{k}$$

A vector  $\vec{F}$  is said to be irrotational if  $\nabla \times \vec{F} = 0$



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

$$\begin{aligned} \text{Now, } \nabla \times \vec{F} &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xy & (x^2 + 2yz) & (y^2 + 1) \end{vmatrix} \\ &= \\ &= \vec{i} \left[ \frac{\partial(y^2 + 1)}{\partial y} - \frac{\partial(x^2 + 2yz)}{\partial z} \right] - \vec{j} \left[ \frac{\partial(y^2 + 1)}{\partial x} - \frac{\partial(2xy)}{\partial z} \right] + \vec{k} \left[ \frac{\partial(x^2 + 2yz)}{\partial x} - \frac{\partial(2xy)}{\partial y} \right] \\ &= \vec{i}(2y - 2y) - \vec{j}(0 - 0) + \vec{k}(2x - 2x) \\ \nabla \times \vec{F} &= \mathbf{0} \end{aligned}$$

**13. Show that the vector  $\vec{F} = 3y^4z^2\vec{i} + 4x^3z^2\vec{j} - 3x^2y^2\vec{k}$  is solenoidal.**

**We know that, if  $\vec{F}$  is solenoidal, we have**

$$\begin{aligned} \text{div } \vec{F} &= \nabla \cdot \vec{F} \\ &= \left( \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \cdot [3y^4z^2\vec{i} + 4x^3z^2\vec{j} - 3x^2y^2\vec{k}] \\ &= \frac{\partial}{\partial x}(3y^4z^2) + \frac{\partial}{\partial y}(4x^3z^2) + \frac{\partial}{\partial z}(-3x^2y^2) \\ &= \mathbf{0+0+0} \\ \therefore \text{div } \vec{F} &= \mathbf{0} \end{aligned}$$

**Hence  $\vec{F}$  is solenoidal.**

**14. Define the line integral.**

Let  $\vec{F}$  be a vector field in space and let AB be a curve described in the sense A to B. Divide the curve AB into n elements  $d\vec{r}_1, d\vec{r}_2, \dots, d\vec{r}_n$ .

Let  $\vec{F}_1, \vec{F}_2, \dots, \vec{F}_n$  be the values of this vector at the junction points of the vectors  $d\vec{r}_1, d\vec{r}_2, \dots, d\vec{r}_n$ , then the sum

$$\lim_{n \rightarrow \infty} \sum_A^B \vec{F}_n \cdot d\vec{r}_n = \int_A^B \vec{F} \cdot d\vec{r} \quad \text{is called the line integral.}$$

**If the line integral is along the curve c then it is denoted by**

$$\int_c \vec{F} \cdot d\vec{r} \quad \text{or} \quad \oint_c \vec{F} \cdot d\vec{r} \quad \text{if } c \text{ is a closed curve.}$$



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

**15. Evaluate  $\int_c \vec{F} \cdot d\vec{r}$  along the curve  $c$  in  $xy$  plane,  $y = x^3$  from the point (1,1) to (2,8) if  $\vec{F} = (5xy - 6x^2)\vec{i} + (2y - 4x)\vec{j}$ .**

**Given**  $\vec{F} = (5xy - 6x^2)\vec{i} + (2y - 4x)\vec{j}$ ,  $y = x^3$

**Now,**  $\vec{r} = x\vec{i} + y\vec{j}$ ;  $d\vec{r} = dx\vec{i} + dy\vec{j}$

**Here**  $y = x^3$ ;  $dy = 3x^2 dx$

$$\begin{aligned} \therefore \int_c \vec{F} \cdot d\vec{r} &= \int_c [(5xy - 6x^2)\vec{i} + (2y - 4x)\vec{j}] \cdot [dx\vec{i} + dy\vec{j}] \\ &= \int_c [(5xy - 6x^2)dx + (2y - 4x)dy] \\ &= \int_c [(5x(x^3) - 6x^2)dx + [(2x^3 - 4x)3x^2 dx]] \\ &= \int_c (5x^4 - 6x^2 + 6x^5 - 12x^3)dx \\ &= x^5 - 2x^3 + x^6 - 3x^4 \end{aligned}$$

**There fore  $\int_c \vec{F} \cdot d\vec{r}$  from the point (1,1) to (2,8)**

**ie.,**  $\int_1^2 \vec{F} \cdot d\vec{r} = [x^5 - 2x^3 + x^6 - 3x^4]_1^2 = 35$

**16. Define surface integral.**

**An integral which is evaluated over a surface is called a surface integral.**

$\therefore \lim_{n \rightarrow \infty} \sum_{i=1}^n \vec{F}(x_i, y_i, z_i) \cdot \hat{n}_i \Delta S_i$  is known as the surface integral.

**17. Find  $\iint_s \vec{r} \cdot d\vec{s}$ , where  $s$  is the surface of the tetrahedron whose**

**vertices are (0,0,0), (1,0,0), (0,1,0), (0,0,1).**

**By Gauss divergence theorem,**

$$\iint_s \vec{r} \cdot d\vec{s} = \iiint_v (\nabla \cdot \vec{r}) dv$$



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---

$$\nabla \cdot \vec{r} = \left( \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right) \cdot [x\vec{i} + y\vec{j} + z\vec{k}] = 1+1+1 = 3$$

$$\therefore \iint_S \vec{r} \cdot \hat{n} ds = \iiint_V 3dv = 3V$$

18. If  $\vec{F} = \text{curl } \vec{A}$ , prove  $\iint_S \vec{F} \cdot \hat{n} ds = 0$ , for any closed surface S.

By Gauss divergence theorem,

$$\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{F} dV = \iiint_V \text{div}(\vec{F}) dV$$

$$= \iiint_V \text{div}(\text{curl } \vec{A}) dV = 0 \quad [\text{since } \text{div}(\text{curl } \vec{A}) = 0]$$

19. Define Volume integral.

An integral which can be evaluated over a volume closed by a surface is called a volume integral. Volume integral can be evaluated by triple integration.

ie.,  $\iiint_V f(x, y, z) dv$

20. State Gauss Divergence theorem.

If  $\vec{F}$  is a vector point function, finite and differentiable in a region  $r$  bounded by a closed surface S, then the surface integral of the normal component of  $\vec{F}$  taken over S is equal to the integral of divergence of  $\vec{F}$  taken over V.

ie.,  $\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{F} dv$  Where  $\hat{n}$  is the unit vector in the positive normal to S.

21. Evaluate  $\iint_S \vec{r} \cdot \hat{n} ds$ , where S is a Closed surface .

By Gauss Divergence theorem , we have

$$\iint_S \vec{r} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{r} dv$$





**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

$$\begin{aligned}
 &= \iiint_V \left[ \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right] (x \vec{i} + y \vec{j} + z \vec{k}) dv \\
 &= \iiint_V \left[ \frac{\partial(x)}{\partial x} + \frac{\partial(y)}{\partial y} + \frac{\partial(z)}{\partial z} \right] dv \\
 &= \iiint_V (1 + 1 + 1) dv = 3 \iiint_V dv = 3V
 \end{aligned}$$

**22.. Prove that**  $\iint_S \phi \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{\phi} dV$

**By Gauss Divergence theorem , we have**  $\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{F} dV$

Let  $F = \phi \vec{c}$  where  $\vec{c}$  is a constant vector. Then ,

$$\iint_S \vec{\phi} \cdot \vec{c} \cdot \hat{n} ds = \iiint_V \nabla \cdot (\phi \vec{c}) dv$$

$$\iint_S \vec{c} \cdot (\vec{\phi} \cdot \hat{n}) ds = \iiint_V \vec{c} \cdot (\nabla \phi) dv$$

**Taking  $\vec{c}$  outside the integrals , we get**

$$\vec{c} \cdot \iint_S \vec{\phi} \cdot \hat{n} ds = \vec{c} \cdot \iiint_V \nabla \phi dv$$

$$\iint_S \vec{\phi} \cdot \hat{n} ds = \iiint_V \nabla \phi dv$$

**23. Evaluate**  $\iint_S xdydz + ydzdx + zxdy$  **over the region of radius a.**

$$\iint_S xdydz + ydzdx + zxdy = \iiint_V \left[ \frac{\partial(x)}{\partial x} + \frac{\partial(y)}{\partial y} + \frac{\partial(z)}{\partial z} \right] dx dy dz$$



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---

$$\begin{aligned}
 &= \iiint_V (1 + 1 + 1) dx dy dz \\
 &= 3 \iiint_V dv = 3v \\
 &= 3 \left[ \frac{4}{3} \pi a^3 \right] = 4\pi a^3
 \end{aligned}$$

**24. State Green's theorem in the plane**

If R is a closed region of the xy-plane bounded by a simple closed curve C and if M and N are continuous derivatives in R, then

$$\int M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy \text{ where C is travelled in the}$$

anti-clockwise direction.

**25. Using Green's theorem, prove that the area enclosed by a simple closed curve C**

is  $\frac{1}{2} \int (x dy - y dx) dx dy$ .

Consider By Green's theorem,

$$\int M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy \dots\dots\dots(1)$$

Consider  $\frac{1}{2} \int (x dy - y dx) dx dy = \int \frac{x}{2} dy - \frac{y}{2} dx = \int -\frac{y}{2} dx + \frac{x}{2} dy$

[since,  $M = -\frac{y}{2}$ ;  $N = \frac{x}{2}$ ]

From (1),  $\int -\frac{y}{2} dx + \frac{x}{2} dy = \iint_R \left[ \frac{1}{2} - \left( -\frac{1}{2} \right) \right] dx dy$

$$= \iint_R dx dy = \text{Area bounded by a closed curve 'C'}$$

**26. State Stoke's theorem.**

If  $\vec{F}$  is any continuous differentiable vector function and S is a surface enclosed by a curve C then,  $\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl } \vec{F} \cdot \hat{n} ds$  where  $\hat{n}$  is the unit normal vector at any point of S.



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

27. Using Stoke's theorem, prove that  $\int_c \vec{r} \cdot d\vec{r} = 0$ .

Given,  $\int_c \vec{r} \cdot d\vec{r}$  where  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$

$$\therefore \int_c \vec{r} \cdot d\vec{r} = \iint_s \text{curl} \vec{r} \cdot \hat{n} \, ds \quad [\because \text{by Stoke's theorem}]$$

$$= 0 \quad \left[ \because \text{curl} \vec{r} = \nabla \times \vec{r} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = 0 \right]$$

28. Find the constants a,b,c so that,  $\vec{F} = (x + 2y + az)\vec{i} + (bx - 3y - z)\vec{j} + (4x + cy + 2z)\vec{k}$  is irrotational.

Given  $\nabla \times \vec{F} = 0$

$$\text{ie.,} \quad \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x + 2y + az & bx - 3y - z & 4x + cy + 2z \end{vmatrix} = 0$$

$$\Rightarrow \vec{i} \left[ \frac{\partial}{\partial y} (4x + cy + 2z) - \frac{\partial}{\partial z} (bx - 3y - z) \right] - \vec{j} \left[ \frac{\partial}{\partial x} (4x + cy + 2z) - \frac{\partial}{\partial z} (x + 2y + az) \right] + \vec{k} \left[ \frac{\partial}{\partial x} (bx - 3y - z) - \frac{\partial}{\partial y} (x + 2y + az) \right] = 0$$

$$\Rightarrow \vec{i}[c + 1] - \vec{j}[4 - a] + \vec{k}[b - 2] = 0\vec{i} + 0\vec{j} + 0\vec{k}$$

$$\Rightarrow c + 1 = 0 \quad 4 - a = 0 \quad b - 2 = 0$$

$$\Rightarrow c = -1 \quad ; \quad a = 4 \quad ; \quad b = 2$$

29. If  $\vec{F} = x^2\vec{i} + xy^2\vec{j}$ , evaluate the line integral  $\int_c \vec{F} \cdot d\vec{r}$  from (0,0) to (1,1)

along the path  $y = x$ .

Given  $\vec{F} = x^2\vec{i} + xy^2\vec{j}$ ,  $x = y$



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

$$dx = dy$$

$$\vec{r} = x\vec{i} + y\vec{j}$$

$$d\vec{r} = dx\vec{i} + dy\vec{j}$$

$$\begin{aligned} \vec{F} \cdot d\vec{r} &= x^2 dx + xy^2 dy = x^2 dx + x^3 dx \quad [ \because x = y, dx = dy ] \\ &= (x^2 + x^3) dx \end{aligned}$$

$$\int_c \vec{F} \cdot d\vec{r} = \int_0^1 (x^2 + x^3) dx = \frac{7}{12}$$

**30. What is the greatest rate of increase of  $\phi = xyz^2$  at  $(1,0,3)$ .**

**Given**  $\phi = xyz^2$

$$\begin{aligned} \nabla\phi &= \vec{i} \frac{\partial\phi}{\partial x} + \vec{j} \frac{\partial\phi}{\partial y} + \vec{k} \frac{\partial\phi}{\partial z} \\ &= \vec{i}(yz^2) + \vec{j}(xz^2) + \vec{k}(2xyz) \end{aligned}$$

$$(\nabla\phi)_{(1,0,3)} = \vec{i}(yz^2) + \vec{j}(xz^2) + \vec{k}(2xyz)$$

**The greatest rate of increase =  $|\nabla\phi| = \sqrt{81} = 9$  units**

**31. Using Green's theorem, find the area of a circle of radius r.**

**We know by Green's theorem,**

$$\text{Area} = \frac{1}{2} \int_c (x dy - y dx)$$

**For a circle of radius r, we have  $x^2 + y^2 = r^2$**

**Put**  $x = r \cos \theta, y = r \sin \theta$

$$dx = -r \sin \theta d\theta, dy = r \cos \theta d\theta \quad [ \theta \text{ varies from } 0 \text{ to } 2\pi ]$$

$$\text{Area} = \frac{1}{2} \int_0^{2\pi} [r \cos \theta r \cos \theta - r \sin \theta (-r \sin \theta)] d\theta$$

$$= \frac{1}{2} \int_0^{2\pi} r^2 d\theta = \frac{1}{2} r^2 [\theta]_0^{2\pi}$$

**Area =  $\pi r^2$  sq.units.**

**32. If  $\nabla\phi$  is solenoidal find  $\nabla^2\phi$ .**



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---

**Given**  $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$  is solenoidal.

$$\therefore \nabla \cdot \nabla \phi = 0$$

**But**  $\nabla^2 \phi = \nabla \cdot \nabla \phi = 0$

**33. If**  $\vec{r} = (x\vec{i} + y\vec{j} + z\vec{k})$ , **find**  $\nabla \times \vec{r}$

**Given**  $\vec{r} = (x\vec{i} + y\vec{j} + z\vec{k})$

$$\nabla \times \vec{r} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \vec{i}(0-0) + \vec{j}(0-0) + \vec{k}(0-0) = \vec{0}$$

**34. Define Volume integral.**

An integral which can be evaluated over a volume closed by a surface is called a volume integral. Volume integral can be evaluated by triple integration. Ie.,  $\iiint_V f(x, y, z) dv$

**35. State Gauss Divergence theorem.**

If  $\vec{F}$  is a vector point function, finite and differentiable in a region  $V$  bounded by a closed surface  $S$ , then the surface integral of the normal component of  $\vec{F}$  taken over  $S$  is equal to the integral of divergence of  $\vec{F}$  taken over  $V$ .

ie.,  $\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{F} dv$  Where  $\hat{n}$  is the unit vector in the positive normal to  $S$ .

**36. Evaluate**  $\iint_S \vec{r} \cdot \hat{n} ds$ , where  $S$  is a Closed surface .

**By Gauss Divergence theorem , we have**

$$\iint_S \vec{r} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{r} dv$$



**SNS COLLEGE OF TECHNOLOGY**  
**Differential Equations And Transforms**  
**VECTOR CALCULUS**

---

$$\begin{aligned}
 &= \iiint_V \left[ \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right] (x \vec{i} + y \vec{j} + z \vec{k}) dv \\
 &= \iiint_V \left[ \frac{\partial(x)}{\partial x} + \frac{\partial(y)}{\partial y} + \frac{\partial(z)}{\partial z} \right] dv \\
 &= \iiint_V (1 + 1 + 1) dv = 3 \iiint_V dv = 3V
 \end{aligned}$$

**37. Prove that  $\iint_S \phi \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{\phi} dV$  By Gauss Divergence theorem , we have**

$$\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V \nabla \cdot \vec{F} dV$$

Let  $F = \phi \vec{c}$  where  $\vec{c}$  is a constant vector. Then ,

$$\begin{aligned}
 \iint_S \vec{\phi} \cdot \hat{n} ds &= \iiint_V \nabla \cdot (\phi \vec{c}) dv \\
 \iint_S \vec{c} \cdot (\hat{\phi} \cdot \hat{n}) ds &= \iiint_V \vec{c} \cdot (\nabla \phi) dv
 \end{aligned}$$

Taking  $\vec{c}$  outside the integrals , we get

$$\begin{aligned}
 \vec{c} \cdot \iint_S \hat{\phi} \cdot \hat{n} ds &= \vec{c} \cdot \iiint_V \nabla \phi dv \\
 \iint_S \hat{\phi} \cdot \hat{n} ds &= \iiint_V \nabla \phi dv
 \end{aligned}$$

**38. Evaluate  $\iint_S xdydz + ydzdx + zdx dy$  over the region of radius a.**

$$\iint_S xdydz + ydzdx + zdx dy = \iiint_V \left[ \frac{\partial(x)}{\partial x} + \frac{\partial(y)}{\partial y} + \frac{\partial(z)}{\partial z} \right] dx dy dz$$



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---

$$\begin{aligned}
 &= \iiint_V (1 + 1 + 1) dx dy dz \\
 &= 3 \iiint_V dv = 3v \\
 &= 3 \left[ \frac{4}{3} \pi a^3 \right] = 4\pi a^3
 \end{aligned}$$

**39. State Green's theorem in the plane**

If  $R$  is a closed region of the  $xy$ -plane bounded by a simple closed curve  $C$  and if  $M$  and  $N$  are continuous derivatives in  $R$ , then

$$\int M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy \quad \text{where } C \text{ is travelled in the anti-}$$

clockwise direction.

**40. Using Green's theorem, prove that the area enclosed by a simple closed curve  $C$**

is  $\frac{1}{2} \int (x dy - y dx) dx dy$ .

consider By Green's theorem,

$$\int M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy \dots\dots\dots(1)$$

Consider  $\frac{1}{2} \int (x dy - y dx) dx dy = \int \frac{x}{2} dy - \frac{y}{2} dx = \int -\frac{y}{2} dx + \frac{x}{2} dy$

[since,  $M = -\frac{y}{2}$ ;  $N = \frac{x}{2}$ ]

From (1),  $\int -\frac{y}{2} dx + \frac{x}{2} dy = \iint_R \left[ \frac{1}{2} - \left( -\frac{1}{2} \right) \right] dx dy$

$$= \iint_R dx dy = \text{Area bounded by a closed curve 'C'}$$

**41. State Stoke's theorem.**



SNS COLLEGE OF TECHNOLOGY  
Differential Equations And Transforms  
VECTOR CALCULUS

---



If  $\vec{F}$  is any continuous differentiable vector function and S is a surface enclosed by a curve C then,  $\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl } \vec{F} \cdot \hat{n} ds$  where  $\hat{n}$  is the unit normal vector at any point of S.

42. If  $\vec{F} = (y^2 \cos x + z^2)\vec{i} + (2y \sin x - 4)\vec{j} + 3xz^2\vec{k}$ , find its scalar potential.

To find  $\phi$  such that  $\vec{F} = \text{grad } \phi$

$$(y^2 \cos x + z^2)\vec{i} + (2y \sin x - 4)\vec{j} + 3xz^2\vec{k} = \vec{i} \frac{\partial \phi}{\partial x} + \vec{j} \frac{\partial \phi}{\partial y} + \vec{k} \frac{\partial \phi}{\partial z}$$

Integrating the equations partially w.r.to x,y,z respectively.

$$\phi = y^2 \sin x + xz^3 + f_1(y, z)$$

$$\phi = y^2 \sin x - 4y + f_2(x, z)$$

$$\phi = xz^3 + f_3(y, z)$$

Therefore  $\phi = y^2 \sin x + xz^3 - 4y + c$  is a scalar potential.